

# F

## Geometric Applications of Matrices

In this Appendix we summarize some geometric applications of matrices. The space is 3-dimensional. Indexed homogeneous Cartesian coordinates<sup>1</sup>  $\{x_0, x_1, x_2, x_3\}$  are used. To pass to physical coordinates, divide  $x_1, x_2$  and  $x_3$  by  $x_0$ :  $\{x_1/x_0, x_2/x_0, x_3/x_0\}$ . If  $x_0 = 0$  the physical coordinates are at infinity.

### §F.1 POINTS, PLANES

Points in 3D space will be identified by  $X, Y, P, Q$ , etc. Their coordinates are put in the 4-vectors

$$\mathbf{x} = [x_0 \ x_1 \ x_2 \ x_3]^T, \quad \mathbf{y} = [y_0 \ y_1 \ y_2 \ y_3]^T, \quad \mathbf{p} = [p_0 \ p_1 \ p_2 \ p_3]^T, \text{ etc} \quad (\text{F.1})$$

Planes in 3D space will be identified by  $A, B, C$ , etc. The equation of plane  $A$  is written  $a_0x_0 + a_1x_1 + a_2x_2 + a_3x_3 = 0$  or

$$\mathbf{a}^T \mathbf{x} = 0, \quad \text{or} \quad \mathbf{x}^T \mathbf{a} = 0 \quad (\text{F.2})$$

The plane  $A$  is thus defined by the 4-vector  $\mathbf{a}$ .

#### EXAMPLE F.1

Find the coordinates of the point where line joining points  $P$  and  $Q$  intersects plane  $A$ .

Solution. Any point of  $PQ$  is  $R$  where

$$\mathbf{r} = \lambda \mathbf{p} + \mu \mathbf{q} \quad (\text{F.3})$$

If  $R$  is on plane  $A$ , then  $\mathbf{a}^T \mathbf{x} = 0$  so that  $\lambda \mathbf{a}^T \mathbf{p} + \mu \mathbf{a}^T \mathbf{q} = 0$ . Absorbing a suitable multiplier into the coordinates of  $R$  we obtain its vector in the form

$$\mathbf{r} = (\mathbf{a}^T \mathbf{q}) \mathbf{p} - (\mathbf{a}^T \mathbf{p}) \mathbf{q}. \quad (\text{F.4})$$

### §F.2 LINES

Let  $\mathbf{x}$  and  $\mathbf{y}$  be the coordinates of points  $X$  and  $Y$  on a given line  $L$ . The  $4 \times 4$  antisymmetric matrix

$$\mathbf{L} = \mathbf{xy}^T - \mathbf{yx}^T \quad (\text{F.5})$$

is called the *coordinate matrix* of the line. It can be shown that  $\mathbf{L}$  determines the line  $L$  to within a scale factor.

Let  $A$  and  $B$  be two planes through  $L$ . The  $4 \times 4$  antisymmetric matrix

$$\mathbf{L}^* = \mathbf{ab}^T - \mathbf{ba}^T \quad (\text{F.6})$$

is called the *dual coordinate matrix* of line  $L$ . It can be shown that<sup>2</sup>

$$\mathbf{L}^* \mathbf{L} = \mathbf{0}. \quad (\text{F.7})$$

---

<sup>1</sup> These coordinates were independently invented in 1827 by Möbius and Feuerbach, and further developed in 1946 by E. A. Maxwell at Cambridge. See E. A. Maxwell, *General Homogeneous Coordinates in Space of Three Dimensions*, Cambridge University Press, 1951.

<sup>2</sup> See E. A. Maxwell, loc. cit., page 150.

**EXAMPLE F.2**

Find where line  $L$  defined by two points  $X$  and  $Y$  meets a plane  $A$ .

Solution. Consider the vector

$$\mathbf{p} = \mathbf{L}\mathbf{a} = (\mathbf{xy}^T - \mathbf{yx}^T)\mathbf{a} = (\mathbf{y}^T\mathbf{a})\mathbf{x} - (\mathbf{x}^T\mathbf{a})\mathbf{y} \quad (\text{F.8})$$

From the last form the point  $P$  of coordinates  $\mathbf{p}$  must lie on the line that joins  $X$  and  $Y$ . Moreover since  $\mathbf{L}$  is antisymmetrical,  $\mathbf{a}^T\mathbf{L}\mathbf{a} = 0$  so that  $\mathbf{a}^T\mathbf{p} = 0$ . Thus  $P$  is the intersection of line  $L$  and plane  $A$ .

**EXAMPLE F.3**

Find the plane  $A$  that joins line  $L$  to a point  $X$ .

Solution.

$$\mathbf{a} = \mathbf{L}^*\mathbf{x} \quad (\text{F.9})$$

where  $\mathbf{L}^*$  is the dual of  $\mathbf{L}$ . The demonstration is trivial.

Two immediate corollaries: (i) Line  $L$  lies in the plane  $A$  if  $\mathbf{L}\mathbf{a} = \mathbf{0}$ ; (ii) Line  $L$  passes through the point  $X$  if  $\mathbf{L}^*\mathbf{x} = \mathbf{0}$ .

**EXAMPLE F.4**

Consider two lines  $L_1$  and  $L_2$  with coordinate matrices  $\mathbf{L}_1, \mathbf{L}_2$  and dual matrices  $\tilde{\mathbf{L}}_1$  and  $\tilde{\mathbf{L}}_2$ , respectively. Find the conditions for the lines to intersect.

Solution. Any of the four equivalent conditions

$$\mathbf{L}_1\mathbf{L}_2^*\mathbf{L}_1 = \mathbf{0}, \quad \mathbf{L}_1^*\mathbf{L}_2\mathbf{L}_1^* = \mathbf{0}, \quad \mathbf{L}_2\mathbf{L}_1^*\mathbf{L}_2 = \mathbf{0}, \quad \mathbf{L}_2^*\mathbf{L}_1\mathbf{L}_2^* = \mathbf{0}. \quad (\text{F.10})$$

For the proof see E. A. Maxwell, loc. cit, page 154.