Coupling from the Atmosphere to Geospace in Antarctica

Xinzhao Chu University of Colorado Boulder CEDAR Prize Lecture 2019 June 18, 2019 @ Santa Fe


Credit: Danny Hampton, Ian Geraghty, and Zimu Li

McMurdo Fe Lidar Observations Since Dec. 2010



McMurdo lidar projects supported by NSF grants OPP-0839091, 1246405, and 1443726


STAR Na Doppler Lidar Added in Jan 2018



By making high-precision laser spectroscopy in space, the neutral temperature, line of sight wind, and Na density are measured simultaneously via detecting the Doppler broadening and bulk Doppler shift of Na D₂ absorption line.

McMurdo lidar projects supported by NSF grants OPP-0839091, 1246405, and 1443726


Simultaneous & Common-Volume Observations with Na Doppler and Fe Boltzmann Lidars at McMurdo





Fe Boltzmann lidar beams & telescopes since Dec 2010

Arrival Heights Lidar Observatory on Ross Island, Antarctica Shooting laser beams at 589, 374 and 372 nm to probe Na and Fe metals, & profile temperatures, vertical winds, and various waves, etc.

Lidar Discovery of Aurora Effect on Fast Amplitude Growth of Temperature Tides in the Thermosphere (Uniqueness of McMurdo: By the Edge of Polar Cap)







(K)

70

60

50

40


30

20

10

Dr. Weichun Fong Winter-over 2013 First Place Prize 2015 CEDAR Students Poster Competition

Lidar-Discovered Thermosphere-Ionosphere Fe Layers (TIFe) Correlated to Solar and Geomagnetic Storms






Dr. Zhibin Yu Winter-over 2011 First Place Prize 2013 CEDAR Students Poster Competition


Courtesy of Dr. Delores Knipp & Dr. Zhonghua Xu Coupling from the Atmosphere to Geospace In Antarctica

> Xinzhao Chu University of Colorado Boulder CEDAR Prize Lecture 2019


Credit: Danny, Ian, and Zimu




Dr. Cao Chen



Arrival Heights is a Hotspot of Gravity Waves



Lidar Discovery of Persistent Gravity Waves with Inertial τ of 3–10 h and λ_z of 20–30 km






Poster Competition


[Chen et al., JGR, 2016]

Persistent, large-amplitudes, dominant in the MLT (T' ~ ±20-30 K) No pause during nearly 3-day observation!!! Occurring on every lidar run; as a group, these waves are perpetual What wave sources could be so persistent??? Non-tidal periods, non-fixed phases, phase traced down to the stratosphere

9


MLT Persistent Gravity Waves in June 2011-2015






Freq-spectral slopes -2.7 below 100 km, gradually become shallower -1.6 at 110 km

Persistent waves aren't tidal waves, aren't atmos. normal modes, unlikely wave-wave interactions, but gravity waves!






Dr. Cao Chen Winter-over 2014




Ian Geraghty Winter-over 2019 Undergrad Honorable Mention 2017

What wave sources could be so persistent???

Sources of the MLT Persistent Waves ??? Traced Back to the Stratosphere




This is the first time inertia-gravity waves (IGWs) observed in the Antarctic MLT by lidar and radar together. [Chen et al., JGR, 2013]





Dr. Adrian McDonald

12




Dr. Jian Zhao Winter-over 2015



Dr. Chihoko Yamashita First Place Prize 2011 CEDAR Students Poster Competition


Statistical Characterization of Dominant Gravity Waves in the Stratosphere (30-50 km)



Т


Dominant GWs in the stratosphere are different from the MLT persistent waves

Stratospheric Gravity Wave Spectra & Characteristics




Dr. Jian Zhao Winter-over 2015

Lognormal distributions of vertical wavelength, period, and vertical phase speed [Zhao et al., JGR, 2017]




GW potential energy density (30-50 km) over 5 years [Chu et al., JGR, 2018]




Epm and N² exhibit seasonal patterns with summer minima & winter maxima

Driving Factors for Epm Seasonal Variations



and Wave Sources [Chu et al., JGR, 2018]






Dr. Lynn Harvey



Stratospheric Epm vs. Polar Vortex Location [Chu et al., JGR, 2018]

Lidar Discoveries Inspired Theoreticians and Modelers to Search for the Wave Sources (Vadas and Becker)




[Becker and Vadas, JGR, 2018; Vadas and Becker, JGR, 2018]





Secondary Gravity Wave Generation by Localized, Intermittent Body Force




[Vadas, Zhao, Chu, and Becker, JGR, 2018]




Dr. Vadas

Theory

A New Picture of Antarctic Gravity Waves in Our Papers



Scientific Merits of the New Understandings




[Chen et al., JGR, 2016]

[Becker and Vadas, JGR, 2018]


TIFe Layers Formed by Plasma-Neutral Coupling

Thermosphere-lonosphere Fe (TIFe) layers [Chu et al., GRL, 2011]






Dr. Zhibin Yu



Dr. Cao Chen

This TIFe layer event on 28 May 2011 demonstrates complex gravity wave activity in Antarctica: 1) 3-10 hr inertial-period gravity waves dominate the temperature variations in the MLT; 2) ~1.5 hr fast gravity waves propagate from the MLT well into the thermosphere.


TIFe Model Simulations and Overall Picture





Dr. Zhibin Yu Winter-over 2011 First Place Prize 2013 CEDAR Students Poster Competition

Forming a Big Picture of Antarctic Gravity Waves




A paradigm shift: Energy and momentum are transferred from lower atmosphere sources to the MLT via a complex multi-step coupling processes involving primary, secondary, and tertiary gravity waves.

Convection is absent from winter Antarctica. Is it possible to form a big picture of gravity wave coupling from near the surface to the thermosphere in Antarctica?



High-Resolution STAR Na Doppler Lidar





@ Arrival Heights


By making high-precision laser spectroscopy in space, the neutral temperature, line of sight wind, and Na density are measured simultaneously via detecting the Doppler broadening and bulk Doppler shift of Na D₂ absorption line.








High-Resolution Vertical Wind & Temp Obs.




High-freq. waves $au \sim 10-20 \text{ min}$ Very obvious on W Still visible on T



- Persistent waves τ ~3-10 hr Dominate T variations Still visible on W
- Resolutions used $\Delta t = 3 - 6 \text{ min}$ $\Delta z = 0.5 - 1 \text{ km}$

High-frequency gravity waves are observed with Na lidar in Antarctica. Both secondary and tertiary gravity wave generation are possible.

Simultaneous TIFe and TINa Observations




8-11 May 2018 Very dynamical TIFe layers with high contrast plus some "regular" TIFe peaking around 6-7 UT

Stunning distinction TIFe vs. TINa Above 105 km

"Diffuse" distribution of TINa throughout night plus TINa layers at time similar to TIFe layers

Mesospheric Fe/Na column abundance ratio is ~3, but the TIFe/TINa ratio varies significantly from <1 to ~55 or higher

High Sensitivity to Detect Diurnal Cycles of TIFe, PMC & V. Winds






Zimu Li Winter-over 2019



Zhuoying Chen

Surprising Results from ~9 Years of Lidar Data

78°S Epm (30-50 km) versus Equatorial QBO Easterly Phase




[Chen, Li, et al., CEDAR poster, 2019]

9 Years of PMC vs. solar cycle or lack of it ?

B_{total} vs Season Mean



Manuel Lindo



2.5

5

10

UT Hour

15

[Lindo et al., CEDAR poster, 2019]

20

25

Concluding Remarks and Outlook

 Synthesis of McMurdo lidar observations, numerical modeling and GW theories leads to a new picture of gravity wave coupling from the lower atmosphere to the thermosphere via secondary GW generation and multi-step coupling concepts. → a paradigm shift?
Still many remaining questions, e.g., wave impacts on transport & circulation, high-freq gravity waves, tertiary waves, MWs, ... questioning our own interpretations every day.
Lidar observations at McMurdo provide huge potentials to the CEDAR--GEM sciences ... What we have studied is just the tip of the iceberg, and many more are awaiting ...



May we use the entire Antarctica as a natural laboratory to advance and test theories of gravity waves, TIMt layers, and A-I-M coupling, etc.?

Gratefully Acknowledge the Tremendous Contributions by Winter-Over Students, Summer Scientists, and Collaborators



TIFe layers: lidar observations and numerical modeling Zhibin Yu PhD 2014 John A. Smith PhD Na and Fe Doppler lidar development & Mach-Zehnder Inter. 2014 Temp tides and aurora effects on temp, & lidar development Weichun Fong PhD 2015 Persistent gravity waves & wave recognition methodology Cao Chen PhD 2016 Jian Zhao Gravity waves in the stratos. & secondary GW generation PhD 2018

Other winter-overs: Zimu Li, Ian Geraghty, Brendan Roberts, Ian Barry, Zhengyu Hua, D. Chang Summer scientists: Wentao Huang, Xian Lu, Zhangjun Wang, Muzhou Lu, Mike Lotto Collaborators: Sharon Vadas, Erich Becker, Chester S. Gardner, Art Richmond, Tim Fuller-Rowell, Lynn Harvey, Adrian McDonald, Mike Jones, John Plane, Jeff Forbes, Bob Robinson, R. Bishop, Zhonghua Xu, Delores Knipp, Hanli Liu, Qian Wu, Mike Taylor, Y. Zhao, D. Pautet, S. Palo,

Special Thanks to Great Mentors



Lidar "Geeks" to Explore the Unknown

Wholehearted Gratitude to NSF, USAP, and Antarctica New Zealand



National Science Foundation -- where discoveries begin Do not follow where the path may lead. Go instead where there is no path & leave a trail.