

1. (34 pt) Evaluate the integral.

(a)  $\int \frac{12}{x^3 + 6x} dx$

(b)  $\int_{1/2}^1 \frac{\sqrt{1-x^2}}{x^2} dx$

(c)  $\int_0^1 \frac{\ln x}{x^2} dx$  (*Hint: first evaluate the indefinite integral*)

2. (20 pt) Consider the region  $\mathcal{R}$  in the first quadrant (Q1) bounded by  $y = x^3 + 1$ ,  $y = 1$ , and  $x = 1$ .

(a) Sketch and shade the region.

(b) Set up (but do not evaluate) integrals to find the following quantities.

i. Volume of the solid generated by rotating  $\mathcal{R}$  about the line  $x = 2$  using the Shell Method

ii. Volume of the solid generated by rotating  $\mathcal{R}$  about the line  $x = 2$  using the Disk/Washer Method

iii. Area of the surface generated by rotating the curve  $y = x^3 + 1$ ,  $0 \leq x \leq 1$ , about the line  $y = 1$

3. (24 pt) Determine if the following expressions converge or diverge. Justify all answers. State the names of any tests or theorems you use.

(a)  $a_n = (-1)^n \frac{\ln(2n)}{\ln(5n)}$

(b)  $\sum_{n=1}^{\infty} \frac{1}{n^2 - \ln n}$

(c)  $\sum_{n=1}^{\infty} \frac{n!}{e^{n^2}}$

4. (8 pt) The  $n$ th partial sum of the series  $\sum_{n=1}^{\infty} a_n$  is  $s_n = \frac{2n}{3n-1}$ .

(a) Find the third term of the series.

(b) Find the sum of the series or explain why it doesn't exist.

5. (20 pt) Be sure to simplify your answers to the following problems.

(a) Evaluate  $\int \cos(\sqrt{x}) dx$  as a power series. (*Hint: Begin with a common Maclaurin series.*)

(b) Find an approximation of  $\int_0^2 \cos(\sqrt{x}) dx$  using the first 2 nonzero terms of the series found in part (a).

(c) Use the Alternating Series Estimation Theorem to find an upper bound for the approximation error. You may assume that the hypotheses of the theorem are satisfied.

6. (18 pt) Consider the parametric curve  $x = 2 \cos t$ ,  $y = 1 + \sin t$  for  $0 \leq t \leq 2\pi$ .

(a) Find a Cartesian equation of the curve. Fully simplify your answer.

(b) Sketch the parametric curve. Indicate with an arrow the direction in which the curve is traced as  $t$  increases.

(c) Find the slope of the line tangent to the curve at  $t = \pi/4$ .

7. (26 pt) Consider the polar curves  $r_1 = \cos(3\theta)$  and  $r_2 = \frac{1}{2}$ .

- (a) Evaluate an integral to find the area of the region in the first quadrant (Q1) inside  $r_1$  and outside  $r_2$ .
- (b) Set up (but do not evaluate) integrals to find the total length of the perimeter (boundary) of the region described in part (a).
- (c) Use the identity  $\cos(3\theta) = 4\cos^3\theta - 3\cos\theta$  to find a Cartesian equation of the curve  $r = \cos(3\theta)$ . It is not necessary to simplify or to solve for  $y$  explicitly.