

1. (34 pt) Evaluate the integral.

(a) $\int \frac{12}{x^3 + 6x} dx$

(b) $\int_{1/2}^1 \frac{\sqrt{1-x^2}}{x^2} dx$

(c) $\int_0^1 \frac{\ln x}{x^2} dx$ (Hint: first evaluate the indefinite integral)

Solution:

(a) (10 pt) Using partial fractions, we have

$$\frac{12}{x^3 + 6x} = \frac{12}{x(x^2 + 6)} = \frac{A}{x} + \frac{Bx + C}{x^2 + 6}.$$

Solving

$$A(x^2 + 6) + x(Bx + C) = 12$$

$$(A + B)x^2 + Cx + 6A = 12$$

gives

$$A + B = 0$$

$$C = 0$$

$$6A = 12$$

which has the solution $A = 2$, $B = -2$, $C = 0$. Therefore

$$\begin{aligned} \int \frac{12}{x^3 + 6x} dx &= \int \left(\frac{2}{x} - \frac{2x}{x^2 + 6} \right) dx \\ &= \boxed{2 \ln |x| - \ln(x^2 + 6) + C} \end{aligned}$$

using the substitution $u = x^2 + 6$, $du = 2x dx$ to integrate $2x/(x^2 + 6)$.

(b) (12 pt) Let $x = \sin \theta$, $dx = \cos \theta d\theta$. Then $\sqrt{1-x^2} = \sqrt{1-\sin^2 \theta} = \cos \theta$ and the new bounds are $\theta = \frac{\pi}{6}$ to $\theta = \frac{\pi}{2}$.

$$\begin{aligned} \int_{1/2}^1 \frac{\sqrt{1-x^2}}{x^2} dx &= \int_{\pi/6}^{\pi/2} \frac{\cos \theta}{\sin^2 \theta} \cdot \cos \theta d\theta = \int_{\pi/6}^{\pi/2} \frac{\cos^2 \theta}{\sin^2 \theta} d\theta \\ &= \int_{\pi/6}^{\pi/2} \frac{1 - \sin^2 \theta}{\sin^2 \theta} d\theta \\ &= \int_{\pi/6}^{\pi/2} (\csc^2 \theta - 1) d\theta \\ &= [-\cot \theta - \theta]_{\pi/6}^{\pi/2} \\ &= \left(0 - \frac{\pi}{2}\right) - \left(-\sqrt{3} - \frac{\pi}{6}\right) = \boxed{\sqrt{3} - \frac{\pi}{3}} \end{aligned}$$

- (c) (12 pt) Apply Integration by Parts with $u = \ln x$, $du = dx/x$, $dv = x^{-2} dx$, $v = -x^{-1}$.

$$\begin{aligned}\int \frac{\ln x}{x^2} dx &= -\frac{\ln x}{x} + \int x^{-2} dx \\ &= -\frac{\ln x}{x} - \frac{1}{x} + C\end{aligned}$$

The definite integral is improper.

$$\begin{aligned}\int_0^1 \frac{\ln x}{x^2} dx &= \lim_{t \rightarrow 0^+} \int_t^1 \frac{\ln x}{x^2} \\ &= \lim_{t \rightarrow 0^+} \left[\frac{-\ln x - 1}{x} \right]_t^1 \\ &= \lim_{t \rightarrow 0^+} \left(-1 + \frac{\ln t + 1}{t} \right) = \boxed{-\infty}\end{aligned}$$

because $\lim_{t \rightarrow 0^+} \ln t = -\infty$ and so $\lim_{t \rightarrow 0^+} \frac{\ln t}{t} = -\infty$. Therefore the integral is divergent.

2. (20 pt) Consider the region \mathcal{R} in the first quadrant (Q1) bounded by $y = x^3 + 1$, $y = 1$, and $x = 1$.


(a) Sketch and shade the region.

(b) Set up (but do not evaluate) integrals to find the following quantities.

- Volume of the solid generated by rotating \mathcal{R} about the line $x = 2$ using the Shell Method
- Volume of the solid generated by rotating \mathcal{R} about the line $x = 2$ using the Disk/Washer Method
- Area of the surface generated by rotating the curve $y = x^3 + 1$, $0 \leq x \leq 1$, about the line $y = 1$

Solution:

(a) (2 pt)



(b) i. (6 pt) $V = \int_a^b 2\pi r h \, dx = \boxed{\int_0^1 2\pi(2-x)x^3 \, dx}$

ii. (6 pt) $V = \int_a^b \pi (R^2 - r^2) \, dx = \boxed{\int_1^2 \pi \left[\left(2 - \sqrt[3]{y-1} \right)^2 - 1 \right] \, dy}$

iii. (6 pt) $S = \int_a^b 2\pi r \sqrt{1 + (y')^2} \, dx = \boxed{\int_0^1 2\pi x^3 \sqrt{1 + (3x^2)^2} \, dx}$

OR $S = \int_a^b 2\pi r \sqrt{1 + (x')^2} \, dy = \boxed{\int_1^2 2\pi(y-1) \sqrt{1 + \left(\frac{1}{3(y-1)^{2/3}} \right)^2} \, dy}$

3. (24 pt) Determine if the following expressions converge or diverge. Justify all answers. State the names of any tests or theorems you use.

(a) $a_n = (-1)^n \frac{\ln(2n)}{\ln(5n)}$

(b) $\sum_{n=1}^{\infty} \frac{1}{n^2 - \ln n}$

(c) $\sum_{n=1}^{\infty} \frac{n!}{e^{n^2}}$

Solution:

(a) (8 pt)

$$\lim_{n \rightarrow \infty} |a_n| = \lim_{n \rightarrow \infty} \frac{\ln(2n)}{\ln(5n)} \stackrel{LH}{=} \frac{\frac{1}{2n} \cdot 2}{\frac{1}{5n} \cdot 5} = \frac{\frac{1}{n}}{\frac{1}{n}} = 1$$

As $n \rightarrow \infty$, the sequence a_n will alternate between values approaching -1 and 1 , so a_n diverges.

(b) (8 pt) Use the Limit Comparison Test and compare to the convergent p-series $\sum_{n=1}^{\infty} \frac{1}{n^2}$.

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{a_n}{b_n} &= \lim_{n \rightarrow \infty} \frac{\frac{1}{n^2 - \ln n}}{\frac{1}{n^2}} = \lim_{n \rightarrow \infty} \frac{1}{n^2 - \ln n} \cdot \frac{1}{\frac{1}{n^2}} \\ &= \lim_{n \rightarrow \infty} \frac{1}{1 - \frac{\ln n}{n^2}} = \frac{1}{1 - 0} = 1 \end{aligned}$$

because $\lim_{n \rightarrow \infty} \frac{\ln n}{n^2} \stackrel{LH}{=} \lim_{n \rightarrow \infty} \frac{\frac{1}{n}}{2n} = 0$.

The limit value is $1 > 0$, so the given series also converges.

(c) (8 pt) Apply the Ratio Test.

$$\begin{aligned} \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| &= \lim_{n \rightarrow \infty} \left| \frac{(n+1)!}{e^{(n+1)^2}} \cdot \frac{e^{n^2}}{n!} \right| = \lim_{n \rightarrow \infty} \frac{(n+1)!}{n!} \cdot \frac{e^{n^2}}{e^{n^2+2n+1}} \\ &= \lim_{n \rightarrow \infty} \frac{n+1}{e^{2n+1}} \stackrel{LH}{=} \lim_{n \rightarrow \infty} \frac{1}{2e^{2n+1}} = 0 < 1 \end{aligned}$$

Therefore the series absolutely converges.

4. (8 pt) The n th partial sum of the series $\sum_{n=1}^{\infty} a_n$ is $s_n = \frac{2n}{3n-1}$.

- (a) Find the third term of the series.
- (b) Find the sum of the series or explain why it doesn't exist.

Solution:

- (a) (4 pt) Because the partial sum $s_2 = a_1 + a_2$ and $s_3 = a_1 + a_2 + a_3$, the third term is

$$a_3 = s_3 - s_2 = \frac{6}{8} - \frac{4}{5} = \boxed{-\frac{1}{20}}.$$

- (b) (4 pt) The sum of the series is

$$\lim_{n \rightarrow \infty} s_n = \lim_{n \rightarrow \infty} \frac{2n}{3n-1} \stackrel{LH}{=} \boxed{\frac{2}{3}}.$$

5. (20 pt) Be sure to simplify your answers to the following problems.

- (a) Evaluate $\int \cos(\sqrt{x}) dx$ as a power series. (*Hint: Begin with a common Maclaurin series.*)
- (b) Find an approximation of $\int_0^2 \cos(\sqrt{x}) dx$ using the first 2 nonzero terms of the series found in part (a).
- (c) Use the Alternating Series Estimation Theorem to find an upper bound for the approximation error. You may assume that the hypotheses of the theorem are satisfied.

Solution:

- (a) (8 pt)

$$\begin{aligned} \cos x &= \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!} \\ \cos \sqrt{x} &= \sum_{n=0}^{\infty} (-1)^n \frac{(\sqrt{x})^{2n}}{(2n)!} = \sum_{n=0}^{\infty} (-1)^n \frac{x^n}{(2n)!} \\ \int \cos \sqrt{x} dx &= \int \sum_{n=0}^{\infty} (-1)^n \frac{x^n}{(2n)!} dx \\ &= \boxed{C + \sum_{n=0}^{\infty} (-1)^n \frac{x^{n+1}}{(2n)!(n+1)}} \end{aligned}$$

(b) (6 pt)

$$\begin{aligned}\int \cos \sqrt{x} dx &\approx \frac{x}{0!1} - \frac{x^2}{2!2} \\ &= x - \frac{x^2}{4} \\ \int_0^2 \cos \sqrt{x} dx &\approx \left[x - \frac{x^2}{4} \right]_0^2 \\ &= \left(2 - \frac{2^2}{4} \right) - 0 = \boxed{1}\end{aligned}$$

(c) (6 pt) Let the partial sum $s_n = \sum_{i=0}^n a_i$. The error when using s_n to estimate the sum S of the series is $|R_n| = |S - s_n|$. By ASET, this error has an upper bound of $|a_{n+1}|$, the next term of the series. For the approximation found in part (b), the error bound is $\frac{2^3}{4!3} = \boxed{\frac{1}{9}}$.

6. (18 pt) Consider the parametric curve $x = 2 \cos t$, $y = 1 + \sin t$ for $0 \leq t \leq 2\pi$.

- (a) Find a Cartesian equation of the curve. Fully simplify your answer.
- (b) Sketch the parametric curve. Indicate with an arrow the direction in which the curve is traced as t increases.
- (c) Find the slope of the line tangent to the curve at $t = \pi/4$.


Solution:

(a) (6 pt)

$$\cos^2 t + \sin^2 t = 1$$

$$\boxed{\frac{x^2}{4} + (y - 1)^2 = 1}$$

(b) (6 pt)



(c) (6 pt)

$$\frac{dy}{dx} = \frac{dy/dt}{dx/dt} = \frac{\cos t}{-2 \sin t}$$


$$\left. \frac{dy}{dx} \right|_{t=\pi/4} = -\frac{\cos \frac{\pi}{4}}{2 \sin \frac{\pi}{4}} = -\frac{\frac{1}{\sqrt{2}}}{2 \cdot \frac{1}{\sqrt{2}}} = \boxed{-\frac{1}{2}}$$

7. (26 pt) Consider the polar curves $r_1 = \cos(3\theta)$ and $r_2 = \frac{1}{2}$.

- Evaluate an integral to find the area of the region in the first quadrant (Q1) inside r_1 and outside r_2 .
- Set up (but do not evaluate) integrals to find the total length of the perimeter (boundary) of the region described in part (a).
- Use the identity $\cos(3\theta) = 4\cos^3\theta - 3\cos\theta$ to find a Cartesian equation of the curve $r = \cos(3\theta)$. It is not necessary to simplify or to solve for y explicitly.

Solution:

(a) (12 pt)



In Q1, the curves intersect at $\cos(3\theta) = \frac{1}{2} \implies 3\theta = \frac{\pi}{3} \implies \theta = \frac{\pi}{9}$.

$$\begin{aligned} A &= \int_{\alpha}^{\beta} \frac{1}{2} (r_1^2 - r_2^2) d\theta \\ &= \int_0^{\pi/9} \frac{1}{2} \left(\cos^2(3\theta) - \frac{1}{4} \right) d\theta \\ &= \int_0^{\pi/9} \frac{1}{2} \cos^2(3\theta) d\theta - \int_0^{\pi/9} \frac{1}{8} d\theta \\ &= \int_0^{\pi/9} \frac{1}{4} (1 + \cos(6\theta)) d\theta - \int_0^{\pi/9} \frac{1}{8} d\theta \\ &= \left[\frac{1}{4} \left(\theta + \frac{1}{6} \sin(6\theta) \right) \right]_0^{\pi/9} - \left[\frac{1}{8} \theta \right]_0^{\pi/9} \\ &= \frac{1}{4} \left(\frac{\pi}{9} + \frac{1}{6} \cdot \frac{\sqrt{3}}{2} \right) - \frac{1}{8} \cdot \frac{\pi}{9} \\ &= \boxed{\frac{\pi}{72} + \frac{\sqrt{3}}{48}} \end{aligned}$$

- (8 pt) The perimeter is formed by r_1 and r_2 for $0 \leq \theta \leq \frac{\pi}{9}$, and a line segment of length $\frac{1}{2}$.

$$\begin{aligned}
L &= \frac{1}{2} + \int_0^{\pi/9} \sqrt{r_1^2 + \left(\frac{dr_1}{d\theta}\right)^2} d\theta + \int_0^{\pi/9} \sqrt{r_2^2 + \left(\frac{dr_2}{d\theta}\right)^2} d\theta \\
&= \boxed{\frac{1}{2} + \int_0^{\pi/9} \sqrt{\cos^2(3\theta) + (-3\sin(3\theta))^2} d\theta + \int_0^{\pi/9} \sqrt{\frac{1}{4}} d\theta} \\
&= \frac{1}{2} + \int_0^{\pi/9} \sqrt{1 + 8\sin^2(3\theta)} d\theta + \frac{\pi}{18}
\end{aligned}$$

(c) (6 pt) Use the given identity and the identities $r^2 = x^2 + y^2$ and $x = r \cos \theta$.

$$r = \cos(3\theta)$$

$$r = 4 \cos^3 \theta - 3 \cos \theta$$

Multiply both sides by r^3 , then substitute.

$$r^4 = 4r^3 \cos^3 \theta - 3r^3 \cos \theta$$

$$(r^2) = 4(r \cos \theta)^3 - 3r^2(r \cos \theta)$$

$$\boxed{(x^2 + y^2)^2 = 4x^3 - 3x(x^2 + y^2)}$$

Alternate Solution: Let $r = \sqrt{x^2 + y^2}$. If $r \neq 0$, then $\cos \theta = x/r$ and

$$r = 4 \cos^3 \theta - 3 \cos \theta$$

$$= 4 \left(\frac{x}{r}\right)^3 - 3 \cdot \frac{x}{r}$$

$$\sqrt{x^2 + y^2} = \frac{4x^3}{(x^2 + y^2)^{3/2}} - \frac{3x}{\sqrt{x^2 + y^2}}.$$