

1. (14 pts) Consider the differential equation

$$x \frac{dy}{dx} = \frac{2x^2 + 1}{e^y}, \quad \text{where } x > 1.$$

Find the particular solution where $y(e) = 0$.

2. (10 pts) A force of 3 pounds will stretch a spring $1/2$ feet beyond its natural length.

- How far will an 8-pound force stretch the spring?
- How much work does it take to stretch the spring to the distance you found in (a)?


On this problem, you may assume that the force F required to stretch a string x units beyond its natural length is given by $F = kx$, where k is a constant.

3. (24 pts) Consider the region $\mathcal{R}$ in Quadrant 1 bounded by $y = \sin(x)$ and $y = \cos(x)$ until their first point of intersection, and the y -axis.

- Sketch the region, and label any intercepts and points of intersection.

For the following parts, set up but DO NOT evaluate integrals for the given quantities:

- the volume of the solid generated by rotating $\mathcal{R}$ around the line $y = -1$;
 - the perimeter of the region;
 - the moment about the x -axis, assuming the region has density ρ .
4. (28 pts) Consider the region below, bounded by the curves $x = 3y^2 - 2$, $x = y^2$, and the x -axis.



- Compute the area of the region.
 - Compute the volume of the solid generated by rotating the region around the line $y = 2$ using the shell method.
 - Consider the portion of the curve $x = y^2$ which bounds the shaded region. Find the surface area of the shape formed by rotating the curve segment around the x -axis.
5. (12 pts) Consider the series given by

$$\frac{1}{a} - \frac{1}{3a^2} + \frac{1}{9a^3} - \frac{1}{27a^4} + \cdots$$

- Write this series in $\sum$ -notation.

- (b) For which values of a will this series converge? Your answer can use inequalities or interval notation.
- (c) Find the sum of the series when $a = \frac{1}{2}$.
6. (12 pts) Determine whether the following quantities converge or diverge. If something converges, determine what it converges to. If something diverges, explain why. *Note: if you use L'Hôpital's Rule, state where you used it. Give the name of any other theorems you use.*

(a) the sequence $a_n = \frac{\ln(n+1)}{\sqrt{n}}$;

(b) the sequence $b_n = \frac{\sin(n) + 1}{n+1}$;

(c) the series $\sum_{n=0}^{\infty} c_n$, which has partial sums given by $s_n = 2 \left(\frac{1}{3} \right)^n$;

(d) the series $\sum d_n$, where the sequence $\{d_n\}$ is given by $d_n = \left(2 - \frac{1}{n} \right)^2$.