

---

INSTRUCTIONS: Outside paper and electronic devices are **not** permitted. Exam is worth 150 points. Neatness counts. Unless indicated, answers with no supporting work may receive no credit. BOX your final answers.

---

1. (40 points) No work is required for number 1, parts (a) thru (t). Simply write an answer and box it:

(a)  $y = \frac{g(x)}{h(x)} \implies \frac{dy}{dx} = ?$

(h)  $f(y) = x \implies \frac{dy}{dx} = ?$

(b)  $y = g(x) \cdot h(x) \implies \frac{dy}{dx} = ?$

(i) What is the volume of a cone?

(c)  $y = [g(x)]^n \implies \frac{dy}{dx} = ?$

(j) What is the volume of a sphere?

(d)  $y = ax^n \implies \frac{dy}{dx} = ?$

(k) What is the surface area of a sphere?

(e)  $y = \csc(x) \implies \frac{dy}{dx} = ?$

(l) Where are the critical points of  $f(x)$  found?

(f)  $y = \cot(x) \implies \frac{dy}{dx} = ?$

(m)  $\sqrt{x^2} = ?$

(g)  $y = \tan(x) \implies \frac{dy}{dx} = ?$

(n)  $\lim_{x \rightarrow 0} \left[ \frac{\sin x}{x} \right] = ?$

(o) What is the limit definition of derivative for  $f(x)$ ?

(p) What must be true of  $f(x)$  in order to apply the Mean Value Theorem?

(q) What must be true of  $f(x)$  in order to apply the Intermediate Value Theorem?

(r) What is the distance between points  $(x_1, y_1)$  and  $(x_2, y_2)$ ?

(s) If  $S(t)$  is the position of an object at time  $t$ , then what is  $S''(t)$ ?

(t) When is  $f(x)$  considered to be continuous at  $x = a$ ?

2. (35 points) Consider the function:  $f(x) = \sqrt{2x+1}$

(a) Find the average rate of change of  $f(x)$  between  $x = 4$  and  $x = \frac{15}{2}$ .

(b) Find the instantaneous rate of change of  $f(x)$  at  $x = 4$ .

(c) Create the linearization of  $f(x)$  at  $a = 4$ .

(d) Use the linearization you created in (c) to estimate  $\sqrt{10}$ .

(e) Is the estimation found in (d) a little too big or a little too small?

(f) We know that  $\sqrt{10} > 3$ . What is the estimated difference?

(g) We know that  $\sqrt{10} > 3$ . What is the true difference? Answer with simplified radicals if needed.

3. (20 points) Consider the function:  $g(x) = \frac{x^4 - 2x^3 + 10x^2 - 20x}{x^2 - 2x}$
- Simplify and reduce  $g(x)$
  - Find the absolute minimum and maximum of  $g(x)$  on the interval  $[3, 5]$ .
  - $\lim_{x \rightarrow \infty} \left[ \frac{1}{g(x)} \right] = ?$
  - Find any horizontal asymptotes of  $h(x) = \frac{x^2}{g(x)} + 2$
4. (20 points) Consider the function  $f(\theta) = \frac{\theta^2 - \theta + \theta \cos^2(\theta)}{-\theta^2}$ .
- Simplify  $f(\theta)$  and write your simplification in terms of  $\sin(\theta)$ .
  - $\lim_{\theta \rightarrow 0} [f(\theta)] = ?$
  - $f' \left( \frac{\pi}{2} \right) = ?$
  - Name two critical points of  $f(\theta)$  found in the interval  $[0, \pi]$ .
5. (20 points) Consider the function:  $f(x) = x^2 - 4x + 7$
- Use the Intermediate Value Theorem to show that  $f(x) = \pi$  for  $2 \leq x \leq 6$ .
  - According to the Mean Value Theorem the average rate of change of  $f(x)$  between  $x = 2$  and  $x = 6$  should equal the instantaneous rate of change of  $f(x)$  for an  $x$ -value inside the interval  $[2, 6]$ . Where does this occur?
  - Name the  $C$ -value guaranteed to exist by Rolle's Theorem for  $f(x)$  on the interval  $[1, 3]$ .
  - Name any vertical asymptotes or holes for  $g(x) = \frac{4x - 4}{f(x) - 4}$
6. (15 points) The following problems are not necessarily related:
- Sketch  $f(x) = \sqrt{(x - 2)^2}$
  - Suppose  $x(x + 2) \leq g(x) + x^2 \leq x^4 + 2$ , then  $\lim_{x \rightarrow 1} [g(x)] = ?$
  - If the radius of a sphere is increasing at  $1/8$  millimeters per second, then how fast is the surface area changing when the diameter is 80 mm?