

INSTRUCTIONS: Books, notes, crib sheets, and electronic devices are not permitted. Write your (1) name, (2) instructor's name, (3) recitation number. Work all problems. Show and explain your work clearly. Note that a correct answer with incorrect or no meaningful supporting work may receive no credit, while an incorrect answer with relevant work may receive partial credit.

- (25 points) Consider the upper half of the sphere described in Cartesian coordinates by $x^2 + y^2 + z^2 \leq 1$. Now, remove the volume of the smaller sphere described in spherical coordinates by $\rho = \cos \phi$.
 - Make a clear sketch of the geometry of the situation as viewed in a plane of constant θ in the cylindrical coordinate system.
 - Set up the integral, in cylindrical coordinates, to determine the volume of material remaining.
 - Set up the integral, in spherical coordinates, to determine the volume of material remaining.
 - Evaluate your integral from either part (b) or (c) above to determine the remaining volume.

- (25 points) Use the transformation $u = x - y$ and $v = 2x + y$ to evaluate the integral

$$I = \iint_R (2x^2 - xy - y^2) dx dy$$

over the finite region R bounded by the lines $y = -2x + 4$, $y = -2x + 7$, $y = x - 2$, and $y = x + 1$.

- Solve for x and y in terms of u and v using the given substitution. Be sure to check this because the rest of the problem depends on this result! Check this again.
- Transform the original region R_{xy} into its corresponding region R_{uv} in the uv -plane. Make a clear sketch of the new region of integration R_{uv} in the uv -plane. Be sure to label all axes, boundaries, intersection points, etc. on your sketch.
- Rewrite the integral for I over the region R_{uv} in the uv -plane in terms of u and v .
- Evaluate I in terms of u and v .

- (25 points) The integral

$$V = \int_{\theta=0}^{2\pi} \int_{\phi=0}^{\pi/2} \int_{\rho=0}^{\cos(\phi)} \rho^2 \sin(\phi) d\rho d\phi d\theta$$

calculates the volume of a spherical object.

- Make a clear sketch of the cross-section of the object in a constant θ plane. (This could be a constant θ plane in either cylindrical coordinates or spherical coordinates.) Clearly label the bounding surfaces of the region of integration.
 - Express V in spherical coordinates using the order $d\phi d\rho d\theta$.
 - Express V in cylindrical coordinates using the order $dr dz d\theta$.
 - Evaluate any of the integrals above (including the original) to determine the volume V .
- (25 points) Consider the counterclockwise circular path in the x - y plane around the origin with a radius of R . Also consider the vector function given by $\mathbf{F} = xy\mathbf{i} + xy\mathbf{j}$.
 - Sketch the entire path in the x - y plane, including the direction of motion. (This should be free points.)
 - Clearly give a parametrization for the path, $\mathbf{r}(t)$, including the limits on t . (Again, this should be free points.)
 - Determine the unit tangent to the path, \mathbf{T} .
 - Determine the **outward** unit normal to the path, \mathbf{n} .
 - Calculate the **flow** along the path C .
 - Calculate the **flux** along the path C .

OVER

Projections and distances $\text{proj}_{\mathbf{A}} \mathbf{B} = \left(\frac{\mathbf{A} \cdot \mathbf{B}}{\mathbf{A} \cdot \mathbf{A}} \right) \mathbf{A}$ $d = \frac{|\overrightarrow{PS} \times \mathbf{v}|}{|\mathbf{v}|}$ $d = \left| \overrightarrow{PS} \cdot \frac{\mathbf{n}}{|\mathbf{n}|} \right|$

Arc length, frenet formulas, and tangential and normal acceleration components

$$ds = |\mathbf{v}| dt \quad \mathbf{T} = \frac{d\mathbf{r}}{ds} = \frac{\mathbf{v}}{|\mathbf{v}|} \quad \mathbf{N} = \frac{d\mathbf{T}/ds}{|d\mathbf{T}/ds|} = \frac{d\mathbf{T}/dt}{|d\mathbf{T}/dt|} \quad \mathbf{B} = \mathbf{T} \times \mathbf{N}$$

$$\frac{d\mathbf{T}}{ds} = \kappa \mathbf{N} \quad \frac{d\mathbf{B}}{ds} = -\tau \mathbf{N} \quad \kappa = \left| \frac{d\mathbf{T}}{ds} \right| = \frac{|\mathbf{v} \times \mathbf{a}|}{|\mathbf{v}|^3} = \frac{|f''(x)|}{|1 + (f'(x))^2|^{3/2}} = \frac{|\dot{x}\ddot{y} - \dot{y}\ddot{x}|}{|\dot{x}^2 + \dot{y}^2|^{3/2}} \quad \tau = -\frac{d\mathbf{B}}{ds} \cdot \mathbf{N}$$

$$\mathbf{a} = a_N \mathbf{N} + a_T \mathbf{T} \quad a_T = \frac{d|\mathbf{v}|}{dt} \quad a_N = \kappa |\mathbf{v}|^2 = \sqrt{|\mathbf{a}|^2 - a_T^2}$$

The Second Derivative Test

Suppose $f(x, y)$ and its first and second partial derivatives are continuous in a disk centered at (a, b) and $f_x(a, b) = f_y(a, b) = 0$. Let $D = f_{xx}f_{yy} - f_{xy}^2$.

1. If $D > 0$ and $f_{xx} < 0$ at (a, b) , then f has a local maximum at (a, b) .
2. If $D > 0$ and $f_{xx} > 0$ at (a, b) , then f has a local minimum at (a, b) .
3. If $D < 0$ at (a, b) , then f has a saddle point at (a, b) .
4. If $D = 0$ at (a, b) , then the test is inconclusive.

Directional derivative, discriminant, and Lagrange multipliers

$$\frac{df}{ds} = (\nabla f) \cdot \mathbf{u} \quad f_{xx}f_{yy} - (f_{xy})^2 \quad \nabla f = \lambda \nabla g, \quad g = 0$$

Taylor's formula (at the point (x_0, y_0))

$$f(x, y) = f(x_0, y_0) + \left[(x - x_0)f_x(x_0, y_0) + (y - y_0)f_y(x_0, y_0) \right]$$

$$+ \frac{1}{2!} \left[(x - x_0)^2 f_{xx}(x_0, y_0) + 2(x - x_0)(y - y_0)f_{xy}(x_0, y_0) + (y - y_0)^2 f_{yy}(x_0, y_0) \right]$$

$$+ \frac{1}{3!} \left[(x - x_0)^3 f_{xxx}(x_0, y_0) + 3(x - x_0)^2(y - y_0)f_{xxy}(x_0, y_0) \right.$$

$$\left. + 3(x - x_0)(y - y_0)^2 f_{xyy}(x_0, y_0) + (y - y_0)^3 f_{yyy}(x_0, y_0) \right] + \dots$$

Linear approximation error

$$|E(x, y)| \leq \frac{1}{2} M (|x - x_0| + |y - y_0|)^2, \quad \text{where } \max\{|f_{xx}|, |f_{xy}|, |f_{yy}|\} \leq M$$

Polar coordinates $x = r \cos \theta \quad y = r \sin \theta \quad r^2 = x^2 + y^2 \quad dA = dx dy = r dr d\theta$

Cylindrical and spherical coordinates

Cylindrical to Rectangular	Spherical to Cylindrical	Spherical to Rectangular
$x = r \cos \theta$	$r = \rho \sin \phi$	$x = \rho \sin \phi \cos \theta$
$y = r \sin \theta$	$z = \rho \cos \phi$	$y = \rho \sin \phi \sin \theta$
$z = z$	$\theta = \theta$	$z = \rho \cos \phi$

$$dV = dx dy dz = r dr d\theta dz = \rho^2 \sin \phi d\rho d\phi d\theta$$

Substitutions in multiple integrals

$$\iint_R f(x, y) dx dy = \iint_S f(x(u, v), y(u, v)) |J(u, v)| du dv \quad \text{where} \quad J(u, v) = \frac{\partial x}{\partial u} \frac{\partial y}{\partial v} - \frac{\partial y}{\partial u} \frac{\partial x}{\partial v}$$

Mass, moments, and center of mass $\text{Mass } M = \iint_R \delta dA$

Moments $M_x = \iint_R y \delta dA \quad M_y = \iint_R x \delta dA$ Center of mass $\bar{x} = M_y/M \quad \bar{y} = M_x/M$

Flow and flux $\text{Flow} = \int_C \mathbf{F} \cdot \mathbf{T} ds = \int_C \mathbf{F} \cdot \mathbf{V} dt = \int_C \mathbf{F} \cdot d\mathbf{r} = \int_C M dx + N dy$

$$\text{Flux} = \int_C \mathbf{F} \cdot \mathbf{n} ds = \int_C M dy - N dx \quad \text{where } \mathbf{n} = \mathbf{T} \times \mathbf{k}$$