
- 1. (24 points) The following problems are unrelated.
  - (a) Find the derivative of  $y = \sqrt{x^5} + \frac{4x^8}{3ax^{2/3}}$  where a is a constant. (Fully simplify your final answer.)
  - (b) Find s'(r) when  $s(r) = \frac{5\cos(r) 8r^2}{(7r^3 + 4)^{102}}$ . (Please DO NOT simplify your final answer.)
  - (c) Consider the function  $h(x) = \sqrt{5-x}$ .
    - i. Determine the linearization, L(x), of y = h(x) at a = 1.
    - ii. Use your linearization from (i) to approximate  $\sqrt{4.1}$ .
- 2. (14 points) Consider the curve given by  $x^2 + 4xy^2 + y^2 = 1$ .
  - (a) Find the value of y' at the point (-4, 1).
  - (b) Determine the normal line to the curve at (-4, 1).
- 3. (15 pts) Consider  $k(x) = x^5(x-3)^7$ .
  - (a) Determine the critical numbers of k(x).
  - (b) For each of the critical numbers you found in (a), determine if it is the location of a "local maximum," "local minimum," or "neither." Clearly state an answer for each critical number, and be sure to justify your answers using a theorem from this class.
- 4. (20 points) Consider  $g(x) = \frac{x}{2x+4}$ .
  - (a) Use the definition of the derivative to show  $g'(x) = \frac{4}{(2x+4)^2}$ . (Note: You must use the definition of the derivative to earn any credit on (a).)
  - (b) Determine the average rate of change of g over [-3, 0].
  - (c) Does g'(x) ever equal the value you obtained in (b)? (Justify your answer.)
  - (d) Why does your answer in (c) not contradict the mean value theorem?
- 5. (15 points) The function f(x) has domain [-1,7]. Consider the graph of y=f'(x) below. The dashed line is the tangent line of y=f'(x) at x=4. Use this graph to answer the questions that follow. No justifications are required on this problem.



(Remember that the graph above is the graph of y = f'(x), not y = f(x).)

- (a) Evaluate  $\lim_{h\to 0} \frac{f'(4+h)-f'(4)}{h}$ .
- (b) On the interval (5,6), is y = f(x) increasing or decreasing?
- (c) On the interval (2,3), is y = f(x) concave up or concave down?
- (d) On which of the following intervals does f(x) have a local extreme value: (-1,1), (3,5) or (5,7)?
- (e) Is the local extreme value noted in (d) a maximum or a minimum?
- (f) If f(4) = 2, what is the equation of the tangent line of y = f(x) at x = 4?
- 6. (12 points) Suppose that two sides of a triangle have fixed lengths of 4 cm and 7 cm, but that the angle between these two sides is growing at a rate of 3 radians per minute. Let  $\theta$  denote the angle between these two sides of the triangle.
  - (a) Draw this triangle and label the angle,  $\theta$ , and the two sides of lengths 4 cm and 7 cm. (We recommend having the side of length 7 cm be the base of your triangle, as this may be helpful in (b).)
  - (b) Determine the rate of change of the area of the triangle with respect to time when the angle is  $\theta = \pi/4$ . (Include the correct units in the final answer.)