

APPM 2350—Section Exam 1—140 points
Wednesday September 26, 6:00pm – 7:30pm, 2018

ON THE FRONT OF YOUR BLUEBOOK write: (1) your name, (2) your student ID number, (3) lecture section/time (4) your instructor's name, and (5) a grading table. Text books, class notes, and calculators are NOT permitted. A one-page one-sided crib sheet is allowed.

Problem 0 – Multiple Choice: (1 point)

Which would you prefer?

- (a) Office hours in ECCR 244 (can be crowded).
- (b) Office hours in FLMG 208 (much larger space, less crowded, farther away from Engineering Center).

Problem 1 – True/False: (25 points)

For the following true/false questions, write TRUE (for always true) or FALSE (if not always true). Your work will not be graded.

- (a) The unit tangent \mathbf{T} , unit normal \mathbf{N} , and unit binormal \mathbf{B} vectors to a space curve $\mathbf{r}(t)$ satisfy the relationships $\mathbf{T} = \mathbf{N} \times \mathbf{B}$, $\mathbf{N} = \mathbf{B} \times \mathbf{T}$, $\mathbf{B} = \mathbf{T} \times \mathbf{N}$.
- (b) An accelerated particle that is moving according to $\mathbf{r}(t) = \langle \cos t^2, \sin t^2 \rangle$, $0 \leq t < \sqrt{2\pi}$ experiences constant curvature.
- (c) A plane is determined by three distinct points that lie on it.
- (d) A line in three dimensions is determined by two distinct points that lie on it.
- (e) The shortest distance between a point P_1 and the line $\mathbf{r}(t)$ defined by the point P_0 and the parallel vector \mathbf{v} with the parametric equation

$$\mathbf{r}(t) = \overrightarrow{OP_0} + t\mathbf{v}, \quad -\infty < t < \infty \quad \text{where } O = (0, 0, 0)$$

is $|\text{proj}_{\mathbf{v}} \overrightarrow{P_0P_1}|$.

Problem 2 – Short Answer Questions: (25 points)

For the questions in this problem, no motivation is required. Clearly box your final answer. Only your boxed final answer will be graded.

- (a) At a certain time t_0 a particle is moving with nonzero speed along a curve in \mathbb{R}^3 such that its velocity and acceleration vectors are scalar multiples of one another. At what rate is the particle's unit tangent vector changing with respect to arc length?
- (b) Consider the collection of vectors of the form $a\mathbf{j} + b\mathbf{k}$ where a and b are real numbers. Find an equation of the plane in which the collection of vectors $\mathbf{i} \times [a\mathbf{j} + b\mathbf{k}]$ lie.
- (c) Describe the surface given by $x^2 - 6x + y^2 - 6y = z^2 + (y - 2)^2 - 13$.
- (d) Suppose the scalar triple product of three vectors \mathbf{u} , \mathbf{v} , \mathbf{w} is $\mathbf{u} \cdot (\mathbf{v} \times \mathbf{w}) = 0$. What conclusion can be drawn about the three vectors?

SEE PROBLEMS ON THE OTHER SIDE

Problem 3: (30 points)

- (a) Show that if the vectors \mathbf{u} and \mathbf{v} have the same length, then $\mathbf{u} + \mathbf{v}$ and $\mathbf{u} - \mathbf{v}$ must be orthogonal.
 (b) You might remember the “law of cosines,” which says that

$$c^2 = a^2 + b^2 - 2ab \cos \theta,$$

where a , b and c are the lengths of the sides of a (not necessarily right) triangle and θ is the angle between the sides of lengths a and b . Use vectors and any properties of vectors learned in class to prove the law of cosines.

- (c) Consider the curve given by $\mathbf{r}(t) = \langle t^2 + 1, t, 0 \rangle$ for $t \geq 0$.
 (i) Find the unit tangent vectors $\mathbf{T}(t)$, $t \geq 0$ and, in particular, $\mathbf{T}(1)$.
 (ii) Sketch a graph of the parametric curve in the plane $z = 0$.
 (iii) **Without calculating** $\mathbf{N}(t)$ (seriously, don't do it), find the unit normal $\mathbf{N}(1)$.
 (iv) Find the unit binormal $\mathbf{B}(1)$.

Problem 4: (30 points)

Consider the lines:

$$\begin{aligned}\mathbf{r}_1(t) &= \langle 5 + 2t, -t, 2 + t \rangle \\ \mathbf{r}_2(t) &= \langle 2 + t, -1 + 2t, -2 + 3t \rangle\end{aligned}$$

- (a) Find the point of intersection of these two lines.
 (b) Find the equation of the plane that contains both of these lines.
 (c) Find the equation of a line that intersects both $\mathbf{r}_1(t)$ and $\mathbf{r}_2(t)$ but is orthogonal to both $\mathbf{r}_1(t)$ and $\mathbf{r}_2(t)$.

Problem 5: (30 points)

A space probe passes through the point $(x, y, z) = (3, -2, 1)$ at time $t = \pi$. Its velocity at time t is given by

$$\mathbf{v}(t) = -3 \cos^2 t \sin t \mathbf{j} + 3 \sin^2 t \cos t \mathbf{k}$$

- (a) Find the location of the space probe when $t = \pi/2$ (give your answer as a point).
 (b) Find the curvature of the space probe's path when $t = \pi/4$.
 (c) Let the point P_0 represent the point in space where the probe is located at time $t = 0$, and P_2 represent the point in space where the probe is located at time $t = \pi/2$. How much further does the probe travel to go from P_0 to P_2 than if it traveled between those points along a straight path?