

APPM 2360: Final Exam

May 6, 2019

ON THE FRONT OF YOUR BLUEBOOK write: (1) your name, (2) your instructor's name, (3) your section number and (4) a grading table. Text books, class notes, cell phones and calculators are NOT permitted. A one page (letter sized **2 sided**) crib sheet is allowed.

Problem 1: (30 points) The following questions are unrelated.

- (a) (12 points) Indicate if the following differential equations are separable or not separable.
- (i) $y' = \frac{t}{y}$
 - (ii) $y' = 6e^{yt}$
 - (iii) $ty' = 2 + y^3$
 - (iv) $y' = \frac{t}{y+1} + \frac{y}{t-1}$
- (b) (18 points) Find the general solution of $ty' = y - t^2$ **using the integrating factor method.**

Problem 2: (50 points; 5 points each) **True/False** (answer True if it is always true, otherwise answer False). No justification is required as there is no partial credit on this question.

- (a) If the row reduced echelon form (RREF) of a matrix A is the matrix R and $|R| = 0$, then $|A| = 0$.
- (b) It is possible for a system of linear differential equations $\mathbf{x}' = \mathbf{A}\mathbf{x}$ to have exactly 2 equilibrium points.
- (c) The solution to the initial value problem given by $\frac{dy}{dt} = \frac{1}{y}$ and $y(0) = 2$ exists and is unique on an interval around $t = 0$.
- (d) The set W of polynomials of degree equal to 2 is a subspace of \mathbb{P}^2 , the space of all polynomials of degree less than or equal to 2.
- (e) In solving $y'' + 4y = \cos(2t)$ using the Method of Undetermined Coefficients, the correct form of the guess for the particular solution is $y_p = A \cos(2t) + B \sin(2t)$.
- (f) The equation $x'' + x' + 9x = 4 \sin(3t)$ has no steady state in the limit $t \rightarrow \infty$.
- (g) Solutions to the forced and damped oscillator equation $y'' + 2y' + 2y = e^{-t} \cos(t)$ will decay to zero in the limit $t \rightarrow \infty$.
- (h) If y_1 and y_2 solve the linear differential equation $L(y) = f(t)$, then $2y_1 - y_2$ also solves $L(y) = f(t)$.
- (i) If $\lambda = 2$ is an eigenvalue of an invertible matrix A , with corresponding eigenvector \mathbf{v} , then $\lambda = 1/2$ is an eigenvalue of the matrix A^{-1} , also with corresponding eigenvector \mathbf{v} .
- (j) A mixture with 1g/L sodium is flowed into a tank at a rate of 1L/min. The resulting mixture flows out of the tank at 1L/min. The tank initially contains 10L of pure water. The amount $x(t)$ in grams of sodium in the tank as a function of time (minutes) is described by the initial value problem:

$$\frac{dx}{dt} = 1 - \frac{x}{10} \quad \text{with} \quad x(0) = 0.$$

Problem 3: (30 points)

- (a) (20 points) You place a pot of boiling water ($T(0) = 100^\circ\text{C}$) outside (where it is $T_a = -20^\circ\text{C}$) at noon; by 1pm it is $T(1) = 20^\circ\text{C}$. Using Newton's Law of Cooling

$$\frac{dT}{dt} = k(T_a - T),$$

where time is in hours, determine how many hours it will be until the water freezes (reaches $T = 0^\circ\text{C}$). Simplify your answer as much as possible.

- (b) (10 points) Consider the following simple model of the zombie apocalypse:

$$\frac{dy}{dt} = y(1 - y),$$

where y is the fraction of the world population who are zombies.

If initially $y(0) = 0$, what fraction of the world y becomes zombies in the limit $t \rightarrow \infty$?

On the other hand if $y(0) = 0.01$, what fraction of the world y becomes zombies as $t \rightarrow \infty$?

Problem 4: (40 points) Consider the following system of differential equations, where we will assume that $x \geq 0$ and $y \geq 0$:

$$\begin{aligned}\frac{dx}{dt} &= x(1 - x - y) \\ \frac{dy}{dt} &= y(2 - x - y)\end{aligned}$$

- (a) (10 points) Find all nullclines and the three equilibrium points with $x \geq 0$ and $y \geq 0$.
 (b) (15 points) Sketch a phase portrait, in the first quadrant ($x \geq 0$ and $y \geq 0$), showing the nullclines, equilibrium points, the direction of the vector field across all the nullclines, and 3 different solution trajectories starting at $(x(0), y(0)) = (2, 3)$; $(x(0), y(0)) = (1, 1/2)$; and $(x(0), y(0)) = (1/4, 1/4)$.
 (c) (15 points) Determine whether each of the three equilibria is stable or unstable.

Problem 5: (30 points)

- (a) (7 points) Determine whether the matrix A is non-singular. If it is non-singular, find its inverse.

$$A = \begin{pmatrix} 3 & 0 & 0 \\ 0 & 6 & 0 \\ 0 & 0 & -2 \end{pmatrix}$$

- (b) (8 points) Find all solutions, \vec{x} , to the problem

$$\begin{aligned}2x_1 - x_2 &= -1 \\ -4x_1 + 2x_2 &= 2.\end{aligned}$$

- (c) (8 points) Do the following vectors form a basis in \mathbb{R}^4 ? Explain why or why not.

$$\begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix}, \quad \begin{pmatrix} 0 \\ 0 \\ 1 \\ 1 \end{pmatrix}, \quad \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \end{pmatrix}, \quad \begin{pmatrix} 0 \\ 1 \\ 1 \\ 1 \end{pmatrix}.$$

- (d) (7 points) Let W be the subspace of \mathbb{R}^3 defined by $W := \{(a, b, c) | a + b + c = 0\}$ find a basis and the dimension of W .

Problem 6: (40 points)

- (a) (10 points) Convert the homogeneous differential equation $y'' + 4y' + 5y = 0$ to a system $\mathbf{x}' = A\mathbf{x}$ of differential equations by setting $x_1 = y$ and $x_2 = y'$, where $A \in \mathbb{R}^{2 \times 2}$.
 (b) (15 points) Find the eigenvalues $\lambda_{1,2}$ and eigenvectors $\mathbf{v}_{1,2}$ of the matrix

$$A = \begin{pmatrix} -1 & -1 \\ 1 & -1 \end{pmatrix}.$$

- (c) (15 points) Write down the general solution \mathbf{x} to $\mathbf{x}' = A\mathbf{x}$ for A defined in part (b).

Problem 7: (30 points) Consider the differential equation $y'' + y = f(t)$ with initial conditions $y(0) = 0$, $y'(0) = 0$, and forcing function $f(t) = 2e^{-t} + \delta(t - c)$.

- (a) (8 points) Compute $F(s)$, the Laplace transform of the forcing function $f(t)$ and state the s values for which it is defined. You may use the Table of Laplace transforms below.
 (b) (8 points) Solve for the Laplace transform of the solution $Y(s)$ to the initial value problem. Again, you may use the Table of Laplace transforms below.
 (c) (14 points) Rearrange $Y(s)$ and invert Laplace transforms to find the solution $y(t)$ to the initial value problem. Simplify the solution as much as possible.

TABLE 1. Table of Laplace transforms

$f(t)$	$F(s)$	s domain	$f(t)$	$F(s)$	s domain
1	$\frac{1}{s}$	$s > 0$	t^n	$\frac{n!}{s^{n+1}}$	$s > 0$, n a positive integer
e^{at}	$\frac{1}{s-a}$	$s > a$	$t^n e^{at}$	$\frac{n!}{(s-a)^{n+1}}$	$s > a$, n a positive integer
$\sin(bt)$	$\frac{b}{s^2 + b^2}$	$s > 0$	$\cos(bt)$	$\frac{s}{s^2 + b^2}$	$s > 0$
$e^{at} \sin(bt)$	$\frac{b}{(s-a)^2 + b^2}$	$s > a$	$e^{at} \cos(bt)$	$\frac{s-a}{(s-a)^2 + b^2}$	$s > a$
$\sinh(at)$	$\frac{a}{s^2 - a^2}$	$s > a $	$\cosh(at)$	$\frac{s}{s^2 - a^2}$	$s > a $
$\delta(t - c)$	e^{-cs}	$c \geq 0, s > 0$	$H(t - c)$	$\frac{e^{-sc}}{s}$	$c \geq 0, s > 0$
$f'(t)$	$sF(s) - f(0)$	depends on $f(t)$	$f(t - c)H(t - c)$	$e^{-cs}F(s)$	$c \geq 0, s > 0$

n^{th} order derivative: $\mathcal{L}\{f^{(n)}\} = s^n \mathcal{L}\{f\} - s^{n-1}f(0) - s^{n-2}f'(0) - \dots - f^{(n-1)}(0)$