

Write your name and your professor's name or your section number in the top right corner of your paper. You are allowed to use textbooks and notes, but you may not ask anyone for help except the professors. To receive full credit on a problem you must show **sufficient justification for your conclusion** unless explicitly stated otherwise. You may use a calculator (or software) to check your answers, but you may not submit decimal answers: you have to show your work.

1. (20 points: 2 each) If the statement is **always true** mark "TRUE"; if it is possible for the statement to be false then mark "FALSE." **No justification is necessary.**

- ____(a) If \mathbf{A} is skew-symmetric and invertible then \mathbf{A}^{-1} is also skew-symmetric.
- ____(b) If $\mathbf{v}_1, \dots, \mathbf{v}_n$ form a basis for \mathbb{R}^n then they also form a basis for \mathbb{C}^n .
- ____(c) Given a basis of \mathbb{R}^n , it is possible to find an inner product such that the basis is orthogonal with respect to the inner product.
- ____(d) If \mathbf{x} is a least-squares solution to $\mathbf{Ax} = \mathbf{b}$, then the residual $\mathbf{b} - \mathbf{Ax}$ is orthogonal to \mathbf{b} . (You may assume orthogonality is defined using the dot product, and the least-squares problem is unweighted.)
- ____(e) If λ_1, \mathbf{v}_1 and λ_2, \mathbf{v}_2 are both eigenvalue/eigenvector pairs for \mathbf{A} , then $\lambda_1\mathbf{v}_1 + \lambda_2\mathbf{v}_2$ is an eigenvector of \mathbf{A} .
- ____(f) If a square matrix \mathbf{A} has the property that $|a_{ii}| > \sum_{j \neq i} |a_{ij}|$ for every i , then \mathbf{A} is invertible.
- ____(g) Suppose that \mathbf{A} is diagonalizable with $\mathbf{A} = \mathbf{S}\mathbf{\Lambda}\mathbf{S}^{-1}$, and that \mathbf{y}^T is a row of \mathbf{S}^{-1} . True or False: $\mathbf{A}^T\mathbf{y} = \lambda\mathbf{y}$ for some eigenvalue λ .
- ____(h) Suppose that \mathbf{A} is a square matrix with real entries and an orthogonal eigenvector basis. True or False: \mathbf{A} is symmetric.
- ____(i) Suppose that \mathbf{Z} is an orthogonal matrix and $\mathbf{A} = \mathbf{Z}\mathbf{B}$ where \mathbf{A} and \mathbf{B} are both real. True or false: \mathbf{A} and \mathbf{B} have the same singular values.
- ____(j) Suppose that \mathbf{A} is 4×10 with rank 4, and that \mathbf{A}^+ is the pseudoinverse of \mathbf{A} . True or false: $\mathbf{AA}^+ = \mathbf{I}$.

2. (16 points) Find the Cholesky decomposition of the following matrix $\mathbf{A} = \begin{bmatrix} 9 & -9 & 9 \\ -9 & 10 & -7 \\ 9 & -7 & 14 \end{bmatrix}$.

3. (20 points) Let

$$L(x, y, z) = \begin{pmatrix} 2(y - z) + x \\ 4z + 2y \\ -z \end{pmatrix}.$$

Find the matrix representation of L with respect to the following basis of \mathbb{R}^3 :

$$\left\{ \begin{pmatrix} 1 \\ -1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 3 \\ 0 \end{pmatrix}, \begin{pmatrix} 2 \\ 0 \\ 1 \end{pmatrix} \right\}.$$

4. (20 points) Suppose that you have the following data from 100 people: weight w_i , height h_i , age t_i and blood pressure p_i (where $i = 1, \dots, 100$). You decide to model the person's blood pressure p as a function of the other factors as follows $p = \beta_0 + \beta_1 \frac{w}{h^2} + \beta_2 t + \beta_3 t^2$. What are the entries of the matrix \mathbf{A} and vector \vec{b} such that the least-squares solution of $\mathbf{A}\vec{x} = \vec{b}$ is the vector of linear regression coefficients β_0, \dots, β_3 ?

5. (24 points) Find the singular value decomposition of $\mathbf{A} = \begin{bmatrix} 1 & 1 \\ -1 & -1 \end{bmatrix}$.