

Topological Colloids & Liquid Crystal Skymions



Ivan I. Smalyukh

*Dept. of Physics, MSE, ECEE, Renewable & Sustainable
Energy Institute, & Soft Materials Research Center
University of Colorado, Boulder*

Boulder, July 17, 2015

Colorado *Physics*
Department of Physics • University of Colorado at Boulder






Thank you !!!








Topology in Different Fields of Science



WORLD SHEET swept out by stringlike particles as they move in spacetime




Green, Scientific American



Optics
 $q=1$





$q=-1$



$q=3$



$q=-3$




Topology and Soft Matter

Intro: Imaging, manipulation, fabrication, & aspects of topology

- Interplay of topologies of surfaces, fields & defects
- 2D “baby” skyrmions & 3D knotted solitons
- Soft matter defects & optical vortices
- Self-assembly & dynamics of knot solitons


Label-free study of the director fields in 3D?

- Polarized self-fluorescence




Multimodal nonlinear optical imaging of LCs

→3D linear/nonlinear optical imaging





→3D optical manipulation



- Broadband CARS-PM
- Broadband SRS-PM
- Multiphoton fluorescence
- Multiple harmonic generation
- Holographic Optical Tweezers

- Label-free chemically-specific orientationally-sensitive 3D imaging
- Simultaneous with non-contact optical manipulation in 3D
- Many imaging modalities with complementary capabilities

Multiphoton self-fluorescence imaging: no dyes




Unpolarized detection:

$$I_{em} \propto I_0 \cos^{2i} \alpha$$

Polarized detection:

$$I_{em} \propto I_0 \cos^{2(i+1)} \alpha$$


5CB



Absorption/emission transition dipole

- Strong orientational sensitivity!
- No dyes needed!!!



Single- vs. Multi-photon Excitation



- probability of simultaneous absorption of two or more photons is very low



- intensity of the fluorescence light falls off very fast above and below focus

- Intrinsic 3D resolution - no pinhole needed**




Coherent anti-Stokes Raman Scattering (CARS)

→ CARS uses two laser frequencies to interact resonantly with a specific molecular vibration;




- 3rd order nonlinear process
- Beat frequency matches that of a Raman vibration -> signal at $\omega_{as} = 2\omega_p - \omega_s$
- Broadband Stokes excitation & CARS detection with Spectral shaping;

Stimulated Raman scattering polarizing m-py



- 2nd order nonlinear process with independent pump/probe and Stokes lasers
- Modulate the Stokes beam – measure the Stimulated Raman Loss signal


Integrated holographic optical tweezers & multimodal 3D imaging: setup schematic



T. Lee, R. P. Trivedi & I. I. Smalyukh, *Opt. Lett.* 35, 3447 (2010);


R.P. Trivedi, T. Lee, K. Bertness, & I.I. Smalyukh, *Opt. Express* 18, 27658-27669 (2010).

Magnetic & optical holonomic control



- Use colloidal objects with super-paramagnetic properties
- Control all degrees of freedom: rotational & translational

Controlling defects through colloidal handles



→ Colloidal dimer of the super-paramagnetic beads


→ What is the defect line?

Varney, Jenness, Smalyukh. Phys Rev E **89**, 022505 (2014)



Simultaneous 3D “drawing” & imaging of defects

3D drawing of defects by a particle

PM Image is not very useful




Slicing the sample





Trivedi, Senyuk, Lee, Smalyukh, *PNAS* **109**, 4744-4749 (2012).

Micro-fabrication of non-spherical colloids




3D complex-shaped colloidal particles

- two-photon photopolymerization





Aqueous Graphene Oxide Colloidal Liquid Crystals

Aqueous GO flakes – 0.3-0.5 vol%, lateral size $\sim 0.4 \mu\text{m}$ on average







→ **Electrostatic stabilization due to the charged groups!**

Textures of a GO nematic liquid crystal at concentration 0.7 wt. %



Defining complex-shaped colloidal microparticles


Optical properties of a GO nematic liquid crystal at concentration 0.7 wt. %




B. Senyuk, N. Behabtu, A. Martinez, T. Lee, D. E. Tsentalovich, G. Ceriotti, J. M. Tour, M. Pasquali, and I. I. Smalyukh. *NATURE COMMUNICATIONS* 6, 7157 (2015)

Discussion/HW Assignment:

how would you fabricate such particles?



Colloidal particles shaped as hollow pyramids



170nm

180nm

270nm

1000nm


1100nm

Tips:

- Note the overall size range provided in nanometers;
- Consider limitations of different techniques;
- Consider approaches not introduced here.


Interplay of topologies of fields, surfaces & defects probed using LC Colloids & Drops

Inclusions/colloids in liquid crystals



Topologically distinct colloidal surfaces?


- Homeomorphic




- Topologically distinct!

Topology of Surfaces: beyond spheres


- Every surface can be assigned an Euler characteristic.
- For a geometry with F faces, V vertices, and E edges, the Euler characteristic is given by $\chi = V - E + F$.
- For an n -torus, the Euler characteristic is given by $\chi = 2 - 2n$.




$$\chi=2$$



$$\chi=2$$



$$\chi=2$$




$$\chi=0$$

Hairy Ball & Poincare-Hopf theorems


Hairy Ball Theorem: no continuous tangent vector field on even-dimensional S^n -spheres (e.g. S^2)

– discussed previously in relation to spherical droplets & particles



Poincare-Hopf theorem: winding numbers of defects in the vector field add to the Euler characteristic: $\chi = 2 - 2g$

$g = 0$



$$\chi = 2$$

$g = 1$



$$\chi = 0$$

$g = 2$



$$\chi = -2$$




$g = 3$



$$\chi = -4$$

...

Defects in 2D: Winding Number




- Director rotates by $2\pi s$ on going around it (s is the winding number)
- Recall this is different in 3D (integer unstable...)

Topology-required surface defects - boojums



Spherical particle with
tangential surface BCs

Two boojums at poles




Winding number of
defects in the 2D field at
the LC particle interface is

$$s = \chi = 2$$



Proc. Natl. Acad. Sci. U.S.A. **103**, 18048 (2006)





Defect-free structures around a ring




Boojums on colloidal rings

Polarizing optical microscopy

3D imaging






2D defect winding numbers







Structural stability

No defects




With boojums



Energetic barrier -
bistable

- Defects are not required but appear to reduce free energy!
- Still satisfy topological constraints (winding numbers add to zero)

Boojums on $g=2$ handlebodies




- Different orientations, but the same net 2D winding numbers of boojums
- Satisfy predictions of topological theorems:

$$\sum_i s_i = \chi = 2 - 2g$$

- To induce boojums of net winding numbers s use a particle of right g

Boojums on colloidal handlebodies




$g=3,4,5$ handlebodies

Theorems & charge conservation:

$$\sum_i s_i = \chi = 2 - 2g$$


g	$\chi = \sum_i s_i$	Number of defects	
		$s = +1$	$s = -1$
1	0	2	2
		0	0
2	-2	2	4
		4	6
3	-4	2	6
		4	8
4	-6	2	8
		4	10
5	-8	2	10
		3	11

Boojums **from handles** on colloidal handlebodies




Strong tangential anchoring promotes boojum splitting into disclination semirings

Splitting of boojum points to semi-rings increases g :
handle-shaped boojums on colloidal handlebodies



Drops of LCs with $g > 0$?


Torus-shaped droplet confinement






- No topological defects induced;

$$\sum_i s_i = \chi = 2 - 2g$$

- Director field reconstructed from 3D 3PEF-PM images






g=2 Torus




$$\sum_i s_i = \chi = 2 - 2g$$

s=-1/2 disclination




Defect Structure

- Rather than boojums, we observe line defects of winding number $-1/2$.
- These structures are topologically equivalent.
- The line defect is lower energy than the boojum in small drops.






$g=3$ torus





- The defects are $-1/2$ line defects spanning across the pore thickness
- They are located at the intersection of the three tori
- The strength at surfaces adds to satisfy topology

$$\sum_i s_i = \chi = 2 - 2g$$





$g=4$ and $g=5$ tori




minimum 6 defects



$$\sum_i s_i = \chi = 2 - 2g$$






minimum 8 defects








Handlebody-shaped colloids, perpendicular BCs


Polarizing
optical m-py



3D imaging





Reconstructed director structures and defects around colloidal handlebodies



Switching director & rotating colloidal tori


- Transformation of defects in applied electric field;
- Bistable particle orientations & fields at no field;



B. Senyuk, Q. Liu, S. He, R. Kamien, R. Kusner, T. Lubensky, & I.I. Smalyukh, " Nature **493**, 200-205 (2013)

Transforming LC defects around colloidal tori

Alignment parallel to the far-field director



Nature **493**, 200 (2013)

- In all configurations bulk defects compensate the charge of particle;
- Satisfy predictions of the Poincare-Hopf index theorem:


$$\sum_i m_i = \pm \chi / 2$$

$$\chi = 2 - 2g$$



- Induce net topological charge $m \pmod{2}$ by a particle of right g

Perpendicular BCs: $g=1$ torus

- Escape in 3D dimension:
 - in cylindrical capillary;
 - under torus confinement;



$$\sum_i m_i = \pm \chi / 2$$



$g=1$ confinement with knotted/linked defect lines

Small drops:








Image comparison:



Links & trefoil knots:





Other torus knots:



Equilibrium structures in g=2 drops

Large drops:




Small drops:




$$\sum_i m_i = \pm \chi / 2$$

Imaging and Elastic Energy


Escaped:




With singular defect lines:





Free energy comparison:






g=4,5 handlebody confinement with “escaped” director field



Polarizing optical m-py:






Nonlinear optical images:






$$\sum_i m_i = \pm \chi / 2$$

Knot-shaped topological colloids (SEM)






<http://georgehart.com/bagel/knot.html>

top view




side view



What defects can be induced by knotted particles?



$T(P,Q)$ knot winds Q times around a circle in interior of a torus & P times around its axis

Euler characteristic of a torus knot




→ Euler characteristic of torus knots is $\chi=0$


Trefoil Colloidal-knot-induced defects



3D image of director structure






Knots induce 12 self-compensating boojums



- 6 $s=1$ & 6 $s=-1$ boojums compensate each other
- consistent with $\chi=0$
- $T(p,q)=T(2,3)$
- The total number of boojums is $4q$


T(3,5) colloidal-knot-induced defects



- 10 $s=1$ & 10 $s=-1$ boojums compensate each other
- consistent with $\chi=0$
- The total number of boojums is $4q$ again



Mutually tangled particle knots & director field

- Trefoil knot colloidal particle;
- perpendicular BCs;
- two knotted $s=-1/2$ defect lines;
- Mutually tangled particle and defect lines!
- Insights into integrations of knotted fields & surfaces




Multi-component particles lacking surface connectivity: colloidal links


Hopf Links




Solomon Links




Colloidal Hopf Links with tangential BCs







Polarizing optical microscopy




Nonlinear depth-resolved optical microscopy




Rotational & translational elastic coupling:





Colloidal Hopf Links with perpendicular BCs






Metastable Hopf links with perpendicular BCs





Rotational & translational elastic coupling:




Colloidal Solomon links with homeotropic BCs



Half-integer line defects follow the particle's tubes



Summary: graph presentations of links



Homeotropic BCs:


- colloidal Hopf links accompanied by 2 or more defect rings
- defect loops have linking number of one or two

- Nonpolar nature provides many ways of satisfying topological constraints of topological interaction between fields, defects & surfaces

2D “baby” skyrmions & 3D
knotted solitons (hopfions),
torons, & umbilics

Links & knots in fields – particles ???



Entire field structure has to conform to the knot!!!




Gaus: Can behave like a particle!!!

- If line crossing is prevented, knots are conserved.

Knot theory motivated by early models of atom



William Thomson, Lord Kelvin



Peter Tait



Periodic Table of the Elements

The periodic table is organized by atomic number (Z) from 1 to 118. Groups are labeled at the top: IA, IBA, IIB, IVB, VB, VIB, VII, IB, and VIIIA. The bottom two rows represent the Lanthanide Series (elements 57-71) and Actinide Series (elements 89-103).

	IA	IIA	III	IVB	V	VI	VII	VIII	IX	X	XI	XII	IIB	IIIA	IVA	VA	VIA	VIIA	VIIIA
1	H													B	C	N	O	F	He
2	Li	Be												Al	Si	P	S	Cl	Ne
3	Na	Mg												Al	Si	P	S	Cl	Ar
4	K	Ca	Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn	Ga	Ge	As	Se	Br	Kr	
5	Rb	Sr	Y	Zr	Nb	Mo	Tc	Ru	Rh	Pd	Ag	Cd	In	Sn	Sb	Te	I	Xe	
6	Cs	Ba	*La	Hf	Ta	W	Re	Os	Ir	Pt	Au	Hg	Tl	Pb	Bi	Po	At	Rn	
7	Fr	Ra	+Ac	Rf	Ha	Sg	Ns	Hs	Mt	110	111	112	113						

* Lanthanide Series
+ Actinide Series

58	59	60	61	62	63	64	65	66	67	68	69	70	71
Ce	Pr	Nd	Pm	Sm	Eu	Gd	Tb	Dy	Ho	Er	Tm	Yb	Lu
90	91	92	93	94	95	96	97	98	99	100	101	102	103
Th	Pa	U	Np	Pu	Am	Cm	Bk	Cf	Es	Fm	Md	No	Lr

Mathematical Theory of Knots & Physics




Edward Witten


- Applications of knot theory in physics;
- Insights into the knot theory from quantum physics & field theories;
- 1990 Fields medal (the only physicist to receive);
- Knotted fields: from elementary particles to cosmology

High dimensional spheres: S^3 in 4D

S^1 in 2 Dimensions



S^2 in 3 Dimensions



S^3 in 4 Dimensions



S^2 - all points in 3D space at distance 1 from the origin.

$$\sqrt{x^2 + y^2 + z^2} = 1$$



S^3 - all points (x,y,z,w) in 4D space at distance 1 from the origin.

$$\sqrt{x^2 + y^2 + z^2 + w^2} = 1$$


High-dimensional spheres?

Stereographic projection

- Stereographic Projection of S^2 in 3D onto a 2D plane
- Stereographic Projection of S^3 in 4D into the 3D space



Stereographic projection in 3D




Circles are linked!

Demo by Paul

2D & 3D solitons: baby Skyrmons vs Hopfions


S^2 in 3 Dimensions



2D Soliton - *baby* Skyrmons in chiral magnets & in chiral liquid crystals


Topology-protected solitons!

S^3 in 4 Dimensions



3D Soliton – Hopfion in chiral LCs (3D Skyrmons)

Mathematical Hopf fibration




Heinz Hopf (1894-1971)


- Part of many theories in mathematics & physics

Nonsingular configurations: Skyrmions, “Baby” Skyrmions...

Skyrmion




Mapping onto
the unit sphere




Skyrmion number - how many times the mapping wraps the unit sphere.


Skyrmion number =2



“Baby Skyrmion” in a confined chiral nematic




$$\pi_2(\mathbb{R}P^2) = \mathbb{Z}$$



- Double-twist cylinder
- Translationally invariant in direction orthogonal to the screen


Hopf Fibration in Molecular Orientation Field



Color-coded molecular orientation field

Pontryagin-Thom construction:

computer-simulated



experimentally reconstructed





Smalyukh et al. Nature Materials 9, 139-145 (2010).
Chen, Ackerman, Alexander, Kamien, Smalyukh Phys. Rev. Lett. **110**, 237801 (2013)

Electrical realignment


- Facile response;
- Field-induced realignment;
- Minimize the electric-field term of free energy density:

$$F_e \propto -(\vec{E} \cdot \hat{n})^2$$



Optical realignment


Optical realignment



→ Realignment by field of laser beam;



$$F_e \propto -(\vec{E} \cdot \hat{n})^2$$

Focused Gaussian beam





$$\text{Intensity} \sim E^2$$

Chiral nematic & vertical boundary conditions



Chiral LC molecule :




→ Vertical BCs incompatible with twist


→ Result – frustration & unwinding

LG beam's effect?

Beam shaping using a spatial light modulator




Optical vortices - phase singularities - screw dislocations




L. Allen et al., PRA 45, 8185 (1992)

•Screw dislocations in the phase of light



J. Courtial et al., Phys Rev Lett 80, 3217, 1998


Control of topological solitons with optical vortices




- Focus 50mW LG beams into a chiral nematic for ~10ms;
- Solitons form spontaneously – different depending on LG charge;
- Stable after turning off the laser light.

3D solitons “Hopfions”: torus knots of nematic field



Double-twist cylinder (recall blue phases)



Double-twist torus




Different (P,Q) -torus knots



→ Realignment by E-field
of vortex laser beams of
different charge;
→ Hopf & Seifert fibrations

Hopf index: linking interpretation


Hopf Fibration in Molecular Orientation Field



Pontryagin-Thom reconstruction:
computer-simulated experimentally reconstructed



Color-coded molecular
orientation field




Hopf index $H=1$




$$\pi_3(\mathbb{R}P^2) = \mathbb{Z}$$

Optically-generated Toron: Hopf + defects




Reconstructed 3D structure



- Focus a 50mW LG beam into the chiral nematic for 10-50ms;
- Toron forms spontaneously;
- Stable after turning off the laser light.

I. Smalyukh *et al.* *Nature Materials* **9**, 139-145 (2010).

Optically-generated Toron: Hopf + defects



- Focus a 50mW LG beam into the chiral nematic for 10-50ms;
- Toron forms spontaneously;
- Stable after turning off the laser light.



I. Smalyukh *et al.* *Nature Materials* **9**, 139-145 (2010).

Twist vs. no twist in a “frustrated” chiral LC




$$f_{elastic} = \frac{K_{11}}{2} (\nabla \cdot \hat{n})^2 + \frac{K_{22}}{2} \left[\hat{n} \cdot (\nabla \times \hat{n}) + \frac{2\pi}{p} \right]^2 + \frac{K_{33}}{2} [\hat{n} \times (\nabla \times \hat{n})]^2$$

Untwisted chiral LC cell




Chiral LC cell with a toron



- Topologically protected: stable unless we induce defects
- Strong energetic barrier $> 1000 K_B T$


Total free energy density isosurfaces



- The energy is minimized in the twisted part of the structure
- Point defects energetically costly


Ferromagnetic Liquid Crystal Colloids

- Platelet-shaped ferromagnetic nanoparticles dispersed in a chiral nematic liquid crystal
- Ferromagnetic nanoplatelets self-align in LC and induce a magnetization field $\mathbf{M}(\mathbf{r})$ parallel $\mathbf{n}(\mathbf{r})$
- Strong sensitivity to weak magnetic field 1-10 G



Free energy, Numerical Modeling

$$F = F_{\text{elast}} + F_{\text{mag}} + F_{\text{coupl}} + F_{\text{surf}}$$




$$F_{\text{elast}} = \frac{1}{2} \int_0^d \left(K_1 (\nabla \cdot \mathbf{n})^2 + K_2 (\mathbf{n} \cdot (\nabla \times \mathbf{n}))^2 + K_3 (\mathbf{n} \times (\nabla \times \mathbf{n}))^2 \right) dx$$

$$F_{\text{mag}} = - \int_0^d \mu_0 \mathbf{M} \cdot \mathbf{H} dx$$

$$F_{\text{coupl}} = - \frac{1}{2} \int_0^d \gamma \mu_0 (\mathbf{n} \cdot \mathbf{M})^2 dx$$


$$F_{\text{surf}} = - \frac{1}{2} W (\mathbf{n} \cdot \mathbf{n}_s)^2$$

Optical Patterning of Director & Magnetization Fields




- Isolated half-integer defect lines allowed in the director field but not in $\mathbf{M}(\mathbf{r})$
- Thus obtain a wall defect in $\mathbf{M}(\mathbf{r})$

Patterning of defects in director & magnetization




→ The wall defect in $\mathbf{M}(\mathbf{r})$ is clearly visible at no applied fields & interacts with domain walls $\mathbf{B}$ is applied

Toron vs. Hopfion: point defects required




- Vector field - 2 oppositely charged point defects
- Help matching the vector field to the far-field $\mathbf{M}$


Topology of magnetization field lines $M(r)$




Toron lateral size vs. field



Toron lateral size vs. field




Comp. simulation: toron lateral size vs. field




Modes of generation: sticking & mobile


At powers $>70\text{mW}$:
→ Pinned where generated;



At powers 30-70mW:
→ Repulsion;
→ Brownian particle-like motion




Dense arrays & crystallization due to confinement & repulsive interactions





Patterning of defects & control of optical solitons

Periodic lattices & quasicrystal-like structures

Honeycomb




Quasicrystal-like pattern






Nature **454**, 501-504;
Ackerman et al, Phys. Rev. E **86**, 021703 (2012)

Example of tuning/erasing toron gratings




Ackerman, Qi, & Smalyukh, *Phys Rev E* (2012)

Diffraction




Stokes Polarimetry for Phase Mapping



$$S_2 = I(\pi / 4, \pi / 4) - I(3\pi / 4, 3\pi / 4) \quad S_3 = I(\pi / 4, 0) - I(3\pi / 4, 0)$$


Phase profile $\delta(x, y) = \arctan[S_3(x, y) / S_2(x, y)]$

Phase singularities generated by toron arrays



P. J. Ackerman, Z. Qi, Y. Lin, C. W. Twombly, M. J. Laviada, Y. Lansac & I. I. Smalyukh, *Scientific Reports* **2**, 414 (2012)





5-7 defects in toron arrays & phase singularities



Self-assembly & non-equilibrium dynamics of knotted 3D solitons


Pair interactions & 2D crystallization

Dense arrays & crystallization




Voltage-reconfigurable structure




vertical cross-section



→ Increasing voltage




→ Structural transformation with umbilical defect formation




Voltage-reconfigurable self-assembly

→ Attractive isotropic @ 2-3V



vertical cross-section





→ Attractive anisotropic dipolar-like @ 3-5V




→ Knowing topology can control self-assembly by U




Field-controlled pair interactions





Field-controlled pair interactions



Voltage-reconfigurable self-assembly



Light-induced dynamics of topological solitons



→ Non-equilibrium dynamics of 2D & 3D solitons

Conclusions

- **Soft Matter** provides an experimental toolkit of exploring topology;
- **Interplay** of defects in light & matter
- **Topology** – the basis for controlling self-assembly

Collaborators

- M. Tasinkevych, MPI, Germany
- R. Kamien & T. Lubensky, UPenn
- R. Kusner, Umass
- J. Yeomans, Oxford
- S. Zumer & M. Ravnik, Ljubljana, Slovenia