

# Antiferromagnetism and Superconductivity

Boulder School for Condensed Matter and Materials Physics  
July 14, 15, 2014  
Subir Sachdev

Talk online: [sachdev.physics.harvard.edu](http://sachdev.physics.harvard.edu)



# Outline


1. Antiferromagnetism and quantum criticality in insulators
2. Onset of antiferromagnetism in metals, and d-wave superconductivity
3. Competing density wave order, and the pseudogap of the cuprate superconductors
4. Non-Fermi liquids

# Outline

1. Antiferromagnetism and quantum criticality in insulators
2. Onset of antiferromagnetism in metals, and d-wave superconductivity
3. Competing density wave order, and the pseudogap of the cuprate superconductors
4. Non-Fermi liquids

# Square lattice antiferromagnet

$$H = \sum_{\langle ij \rangle} J_{ij} \vec{S}_i \cdot \vec{S}_j$$



Ground state has long-range Néel order


Order parameter is a single vector field  $\vec{\varphi} = \eta_i \vec{S}_i$

$\eta_i = \pm 1$  on two sublattices

$\langle \vec{\varphi} \rangle \neq 0$  in Néel state.

# Square lattice antiferromagnet


$$H = \sum_{\langle ij \rangle} J_{ij} \vec{S}_i \cdot \vec{S}_j$$



Weaken some bonds to induce spin entanglement in a new quantum phase



# Square lattice antiferromagnet

$$H = \sum_{\langle ij \rangle} J_{ij} \vec{S}_i \cdot \vec{S}_j$$




Ground state is a “quantum paramagnet”  
with spins locked in valence bond singlets

$$\text{[Diagram of a valence bond singlet]} = \frac{1}{\sqrt{2}} \left( |\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle \right)$$




$$= \frac{1}{\sqrt{2}} (|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle)$$




Quantum critical point with non-local entanglement in spin wavefunction


# Excitation spectrum in the paramagnetic phase




# Excitation spectrum in the paramagnetic phase




# Excitation spectrum in the paramagnetic phase



# Excitation spectrum in the paramagnetic phase




# Excitation spectrum in the paramagnetic phase




Sharp spin 1  
particle excitation  
above an energy  
gap (spin gap)

# Excitation spectrum in the Néel phase




# Excitation spectrum in the Néel phase



Spin waves


# Excitation spectrum in the Néel phase



Spin waves

Derivation of  
field theory of  
critical point


# Description using Landau-Ginzburg field theory



$O(3)$  order parameter  $\vec{\varphi}$


$$\mathcal{S} = \int d^2r d\tau \left[ (\partial_\tau \varphi)^2 + c^2 (\nabla_r \vec{\varphi})^2 + (\lambda - \lambda_c) \vec{\varphi}^2 + u (\vec{\varphi}^2)^2 \right]$$

# Excitation spectrum in the paramagnetic phase




$$V(\vec{\varphi}) = (\lambda - \lambda_c) \vec{\varphi}^2 + u (\vec{\varphi}^2)^2$$

$\lambda > \lambda_c$




# Excitation spectrum in the paramagnetic phase




$$V(\vec{\varphi}) = (\lambda - \lambda_c) \vec{\varphi}^2 + u (\vec{\varphi}^2)^2$$

$\lambda > \lambda_c$




# Excitation spectrum in the paramagnetic phase




$$V(\vec{\varphi}) = (\lambda - \lambda_c) \vec{\varphi}^2 + u (\vec{\varphi}^2)^2$$

$\lambda > \lambda_c$




# Excitation spectrum in the paramagnetic phase




$$V(\vec{\varphi}) = (\lambda - \lambda_c) \vec{\varphi}^2 + u (\vec{\varphi}^2)^2$$

$\lambda > \lambda_c$




# Excitation spectrum in the paramagnetic phase





$$V(\vec{\varphi}) = (\lambda - \lambda_c) \vec{\varphi}^2 + u (\vec{\varphi}^2)^2$$

$\lambda > \lambda_c$




# Excitation spectrum in the Néel phase





$$V(\vec{\varphi}) = (\lambda - \lambda_c)\vec{\varphi}^2 + u(\vec{\varphi}^2)^2$$

$$\lambda < \lambda_c$$




Spin waves (“Goldstone” modes)  
and a longitudinal “Higgs” particle

# Excitation spectrum in the Néel phase





$$V(\vec{\varphi}) = (\lambda - \lambda_c)\vec{\varphi}^2 + u(\vec{\varphi}^2)^2$$

$$\lambda < \lambda_c$$




Spin waves (“Goldstone” modes)  
and a longitudinal “Higgs” particle

# Excitation spectrum in the Néel phase




$$V(\vec{\varphi}) = (\lambda - \lambda_c)\vec{\varphi}^2 + u(\vec{\varphi}^2)^2$$


$$\lambda < \lambda_c$$



Spin waves (“Goldstone” modes)  
and a longitudinal “Higgs” particle



$$\begin{aligned}
 & \text{Diagram of two blue dots in a blue oval} \\
 & = \frac{1}{\sqrt{2}} (|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle)
 \end{aligned}$$



$O(3)$  order parameter  $\vec{\varphi}$

CFT3

$$\mathcal{S} = \int d^2r d\tau \left[ (\partial_\tau \varphi)^2 + c^2 (\nabla_r \vec{\varphi})^2 + s \vec{\varphi}^2 + u (\vec{\varphi}^2)^2 \right]$$

# Quantum Monte Carlo - critical exponents

Table IV: Fit results for the critical exponents  $\nu$ ,  $\beta/\nu$ , and  $\eta$ . We summarize results including a variation of the critical point within its error bar. For the ladder model (top group of values) fit results and quality of fits are also given at the previous best estimate of  $\alpha_c$ . The bottom group are results for the plaquette model. Numbers in [...] brackets denote the  $\chi^2/\text{d.o.f.}$  For comparison relevant reference values for the 3D  $O(3)$  universality class are given in the last line.

$\alpha_c$	$\nu^a$	$\beta/\nu^b$	$\eta^c$
1.9096 $-\sigma$	0.712(4) [1.8]	0.516(2) [0.5]	0.026(2) [0.2]
1.9096	0.711(4) [1.8]	0.518(2) [1.1]	0.029(5) [0.8]
1.9096 $+\sigma$	0.710(4) [1.8]	0.519(3) [2.5]	0.032(7) [1.4]
1.9107 <sup>d</sup>	0.709(3) [1.7]	0.525(8) [15.3]	0.051(10) [12]
1.8230 $-\sigma$	0.708(4) [0.99]	0.515(2) [0.84]	0.025(4) [0.15]
1.8230	0.706(4) [1.04]	0.516(2) [0.40]	0.028(3) [0.31]
1.8230 $+\sigma$	0.706(4) [1.10]	0.517(2) [1.6]	0.031(5) [0.80]
Ref. 49	0.7112(5)	0.518(1)	0.0375(5)

<sup>a</sup> $L > 12$ .

<sup>b</sup> $L > 16$ .

<sup>c</sup> $L > 20$ .

<sup>d</sup>Previous best estimate of Ref. 19.

S. Wenzel and W. Janke, *Phys. Rev. B* **79**, 014410 (2009)

M. Troyer, M. Imada, and K. Ueda, *J. Phys. Soc. Japan* (1997)

# Quantum Monte Carlo - critical exponents

Table IV: Fit results for the critical exponents  $\nu$ ,  $\beta/\nu$ , and  $\eta$ . We summarize results including a variation of the critical point within its error bar. For the ladder model (top group of values) fit results and quality of fits are also given at the previous best estimate of  $\alpha_c$ . The bottom group are results for the plaquette model. Numbers in [...] brackets denote the  $\chi^2/\text{d.o.f.}$  For comparison relevant reference values for the 3D  $O(3)$  universality class are given in the last line.

$\alpha_c$	$\nu^a$	$\beta/\nu^b$	$\eta^c$
1.9096 $-\sigma$	0.712(4) [1.8]	0.516(2) [0.5]	0.026(2) [0.2]
1.9096	0.711(4) [1.8]	0.518(2) [1.1]	0.029(5) [0.8]
1.9096 $+\sigma$	0.710(4) [1.8]	0.519(3) [2.5]	0.032(7) [1.4]
1.9107 <sup>d</sup>	0.709(3) [1.7]	0.525(8) [15.3]	0.051(10) [12]
1.8230 $-\sigma$	0.708(4) [0.99]	0.515(2) [0.84]	0.025(4) [0.15]
1.8230	0.706(4) [1.04]	0.516(2) [0.40]	0.028(3) [0.31]
1.8230 $+\sigma$	0.706(4) [1.10]	0.517(2) [1.6]	0.031(5) [0.80]
Ref. 49	0.7112(5)	0.518(1)	0.0375(5)

Field-theoretic  
RG of CFT3  
E.Vicari *et al.*

<sup>a</sup> $L > 12$ .

<sup>b</sup> $L > 16$ .



<sup>c</sup> $L > 20$ .

<sup>d</sup>Previous best estimate of Ref. 19.



S. Wenzel and W. Janke, *Phys. Rev. B* **79**, 014410 (2009)

M. Troyer, M. Imada, and K. Ueda, *J. Phys. Soc. Japan* (1997)


# TlCuCl<sub>3</sub>



# TlCuCl<sub>3</sub>



An insulator whose spin susceptibility vanishes exponentially as the temperature  $T$  tends to zero.



← Pressure in  $\text{TlCuCl}_3$

# TlCuCl<sub>3</sub> at ambient pressure





FIG. 1. Measured neutron profiles in the  $a^*c^*$  plane of TlCuCl<sub>3</sub> for  $i = (1.35, 0, 0)$ ,  $ii = (0, 0, 3.15)$  [r.l.u.]. The spectrum at  $T = 1.5 \text{ K}$


N. Cavadini, G. Heigold, W. Henggeler, A. Furrer, H.-U. Güdel, K. Krämer and H. Mutka, *Phys. Rev. B* 63 172414 (2001).

# TlCuCl<sub>3</sub> at ambient pressure



N. Cavadini, G. Heigold, W. Henggeler, A. Furrer, H.-U. Güdel, K. Krämer and H. Mutka, *Phys. Rev. B* 63 172414 (2001).

# TiCuCl<sub>3</sub> with varying pressure



Observation of 3 → 2 low energy modes, emergence of new longitudinal mode (the “Higgs boson”) in Néel phase, and vanishing of Néel temperature at quantum critical point

Christian Ruegg, Bruce Normand, Masashige Matsumoto, Albert Furrer, Desmond McMorrow, Karl Kramer, Hans-Ulrich Gudel, Severian Gvasaliya, Hannu Mutka, and Martin Boehm, *Phys. Rev. Lett.* **100**, 205701 (2008)

# Prediction of quantum field theory


Potential for  $\vec{\varphi}$  fluctuations:  $V(\vec{\varphi}) = (\lambda - \lambda_c)\vec{\varphi}^2 + u(\vec{\varphi}^2)^2$

Paramagnetic phase,  $\lambda > \lambda_c$

Expand about  $\vec{\varphi} = 0$ :

$$V(\vec{\varphi}) \approx (\lambda - \lambda_c)\vec{\varphi}^2$$

Yields 3 particles with energy gap  $\sim \sqrt{(\lambda - \lambda_c)}$



# Prediction of quantum field theory

Potential for  $\vec{\varphi}$  fluctuations:  $V(\vec{\varphi}) = (\lambda - \lambda_c)\vec{\varphi}^2 + u(\vec{\varphi}^2)^2$


Paramagnetic phase,  $\lambda > \lambda_c$

Expand about  $\vec{\varphi} = 0$ :

$$V(\vec{\varphi}) \approx (\lambda - \lambda_c)\vec{\varphi}^2$$

Yields 3 particles with energy gap  $\sim \sqrt{(\lambda - \lambda_c)}$

---




Néel phase,  $\lambda < \lambda_c$

Expand  $\vec{\varphi} = (0, 0, \sqrt{(\lambda_c - \lambda)/(2u)}) + \vec{\varphi}_1$ :

$$V(\vec{\varphi}) \approx 2(\lambda_c - \lambda)\varphi_{1z}^2$$


Yields 2 gapless spin waves and one Higgs particle with energy gap  $\sim \sqrt{2(\lambda_c - \lambda)}$



# Prediction of quantum field theory

$$\frac{\text{Energy of Higgs particle}}{\text{Energy of triplon}} = \sqrt{2}$$

$$V(\vec{\varphi}) = (\lambda - \lambda_c)\vec{\varphi}^2 + u(\vec{\varphi}^2)^2$$



Quantum  
criticality at  
non-zero  
temperature