

Post Exposure Plan for Shigella flexneri

Background Information:

Shigella flexneri is found worldwide and is most common in developing countries with poor hygiene and limited clean drinking water. *S. flexneri* causes acute intestinal infections ranging from mild diarrhea to shigellosis (an infection characterized by high fever, abdominal pain, bloody diarrhea with mucous, nausea and vomiting. Stools can be frequent and of small volumes). Complication from *S. flexneri* can include moderate dehydration, hyponatremia (low sodium), seizures (pediatric cases), intestinal complications, bacteremia, and reactive arthritis after the infection.

The primary host of *S. flexneri* is humans with some infections observed in higher primates.

The most common modes of transmission are ingesting contaminated foods or water and the fecal-to-oral route. This organism can survive for months on dry surfaces, increasing chances of exposure. Only a small number of Shigella bacteria is required for an infection.

Shigella species of bacteria are the most frequently identified cause of laboratory-acquired infections because of their high virulence and low infectious dose. Laboratory-acquired infections are presumed to occur via the oral route or eating with contaminated hands.

Shigellosis is a reportable illness. The State of Colorado requires reporting from both physicians and laboratories all instances of infection with Shigella species.

Primary hazards in the laboratory :

Creation of splashes or aerosols, exposure to mucous membranes or skin, ingestion, and exposure to contaminated sharps.

Acceptable disinfectants:

S. flexneri is susceptible to 70% ethanol, 1% sodium hypochlorite, 2% glutaraldehyde, iodines, phenolics and formaldehyde.

Exposure controls and personal protection:

S. flexneri is a Risk Group 2 organism. BSL-2 practices, containment equipment, and facilities are required when working with materials known or suspected of containing this agent.

Personal protective equipment such as lab coats and gloves must be worn when handling infected or potentially infected materials. Eye protection must be used when there is a known or potential risk of generating splashed or aerosols.

Procedures that may generate aerosols or splashes should be conducted in a certified biosafety cabinet.



Personnel should wash their hands frequently while working in and before leaving the laboratory and especially before consuming food. Personal items including waters bottles, cell phones, car keys, etc., should be stored in such a way as to avoid contamination and should be accessed only when the laboratory exercise is completed, lab coats and gloves are removed, and hands are washed.

The Principal Investigator (PI), or their designee, must ensure that all personnel are adequately trained in safe laboratory practices, universal precautions, and proper surface and equipment disinfection before initiating any work with this agent. The PI must also ensure that all personnel are aware of the signs and symptoms of a potential infection with *S. flexneri* with an emphasis on the effects on the immunocompromised.

Researchers who work with this organism should know the antibiotic sensitivity of their study species and whether their species carries the gene responsible for Hemolytic Uremic Syndrome (HUS). In the event of a laboratory-acquired infection, treating health care providers should be informed, so treatment and monitoring are appropriate.

At risk populations:

Immunocompromised individuals are at highest risk for *S. flexneri* infections. These may also include persons with diabetes and those who take medications to reduce their stomach acidity.

Immediate Action by Route of Exposure:

Needlestick, Animal Bite, Laceration: Wash area thoroughly with soap and running water. Do not apply disinfectant to the skin.

Mucous membranes (Eyes, nose, mouth): Flush the eyes for 10-15 minutes if eyes have been exposed to splash or spray containing bacteria. Rinse out mouth without swallowing after any exposure. **Inhalation:** If contaminated materials are aerosolized and potentially inhaled, rinse mouth twice and spit. Do not swallow.

Intact skin: Wash the area thoroughly with soap and water.

After First Aid:

Suspected infections related to research exposures must be reported to the University of Colorado, Boulder's Biosafety Officer at 303-492-2817. Please see below for complete instructions related to suspected research exposure. Treating medical providers should be informed of the possibility of *S. flexneri* infection. The incubation period for this bacteria ranges from 1-7 days.

Post-exposure Prophylaxis:

None.

Symptoms of infection in adults:

Mild infection:



• Diarrhea, which may be self-limiting

Shigellosis:

- Abdominal cramps and tenesmus (feeling that you have to evacuate your gut)
- Frequent low volume stools
- Bloody stools with mucous
- Fever (sometimes > 39°C or >102°F)
- Nausea and vomiting

After First Aid – Treatment and Reporting:

University of Colorado Boulder: Procedures for Work-Related Injuries or Illness, Including Animal Bites, Severe Allergic Symptoms, and Sharps Exposures.

It is the policy of the University of Colorado at Boulder (UCB) that all incidents that result in an injury or severe illness to faculty, staff or students be appropriately documented and reported. If a work-related incident, accident, injury or illness occurs:

a) Medical Treatment

- a. In case of life or limb-threatening emergency call 911 or go immediately to the nearest emergent or urgent care facility. Immediately administer appropriate first aid, including thoroughly washing any wounds or exposed areas with soap and water, if at all possible.
- b. If you are an employee of the University and you need non-emergency care for a work-related illness or injury that has occurred during regular weekday working hours, you must be treated at one of the UCB Designated Medical Providers (DMPs: shown below).
 Boulder Community Hospital Emergency Department has the ability to rapidly test for work-related infectious disease or potential biological exposures; all other DMPs may not have this rapid capability. Testing can be done on the exposed personnel; it may be difficult to test any source patient or sample at the DMP.
- c. After hours or while traveling, go to the nearest urgent or emergent care facility.
- d. **Sharps injuries** that include exposure to human blood, body fluids, tissues, tissue culture cells are considered to be injuries with a potential for transmitting bloodborne pathogens. Prompt evaluation and treatment is necessary for these injuries. (go to Boulder Community Hospital Emergency Department)

b) Reporting

- Report the work-related injury or illness to your supervisor immediately. You or your supervisor should notify the Biosafety Officer at 303-492-2817 or at <u>Cher.Masini@colorado.edu</u>. (The BSO may confer with IACUC administrator, veterinarian, or Occupational Health RN.)
- b. You must file a worker's compensation injury report form **within 4 days of the workrelated injury** / exposure or illness onset. Report the incident on the URM website and use the on-line reporting form.
- c. Sharps injuries must be reported on the URM's needle stick exposure report form.



- d. <u>All injury reporting forms can be found at the URM's website at https://www.cu.edu/risk/file-claim</u>
- c) Eligibility (who is eligible to be seen by UCB Designated Medical Providers?)
 - a. All UCB employees, paid UCB staff, graduate students receiving a traineeship or stipend administered by UCB, undergraduate work-study students and paid undergraduate student assistants are eligible to be seen by UCB DMPs. If you are an UCB employee and you visit your regular primary care provider for a work-related injury and your visit is coded as a work-related injury, your primary insurance may not cover the cost of your visit or treatment with your primary care provider.

Some individuals are not covered by UCB Worker's Compensation: All visiting or resident scholars who do not receive payment via UCB (e.g., Howard Hughes Medical Institute Fellows) must follow the work-related exposure / illness or injury protocol outlined by their parent institution or outside funding source. Contract or consulting employees are also not covered by UCB Worker's Compensation; they need to follow the work-related exposure / illness or injury protocols established by their parent institution or consult with their personal health care provider. Volunteers, contract employees and students not paid by UCB are **not** covered by UCB Worker's Compensation and should be seen by their personal health care provider. If you are a student with a Wardenburg Health care plan, post-exposure lab tests are available and covered.

d) Payment and questions:

- All bills from medical providers must be sent to University Risk Management: University Risk Management (<u>https://www.cu.edu/risk/file-claim</u>) 1800 Grant Street, Ste 700 Denver, CO 80203 Fax: 303-860-5680
- b. For further questions, contact URM at: 303-860-5682 or 888-812-9601

UCB Designated Medical Providers

This list changes frequently. For a current listing of DMPs, please refer to the University of Colorado's Risk Management website at: <u>https://www.cu.edu/risk/dmp</u>