



(A) Biofilm development on indwelling medical device (Image Credit: CanstockPhoto) (B) Biofilm formation seen on the international space station (Image Credit: NASA)

- Silicone is a commonly used substrate in both hospital and spaceflight environments
- With the growing push for extraterrestrial colonization and long-term spaceflight, there exists a need to identify whether similar changes in morphology occur in varying gravitational environments and whether these changes lend themselves to altered virulence.

OBJECTIVES

OBJECTIVE 1. Demonstrate planktonic *P. aeruginosa* growth in varying gravitational regimes via colony forming units (CFU) and biomass growth curves over time.

OBJECTIVE 2. Demonstrate *P. aeruginosa* biofilm growth on silicone at varying gravitational regimes via biomass, mean thickness, substratum coverage, and roughness.

Characterization of *P. aeruginosa* Growth Patterns Under Varying Gravitational Conditions

¹ Modern Human Anatomy Program, University of Colorado School of Medicine, ² BioServe Space Technologies, University of Colorado Boulder,





2.12 1.94 0.95 0.95

Biomass (µm^3/µm^2)

Mean Thickness un

Planktonic *P. aeruginosa* demonstrated significantly different (p<0.01) (A) final cell counts and (B) biomass values between each gravitational regime, with µg demonstrating the greatest values and 1g demonstrating the lowest values. (C) Biofilm morphometrics demonstrated clear patterns of growth with decreasing gravitational regime. Biomass, thickness, and substratum coverage followed an increasing trend with decreasing gravity and biofilm roughness demonstrated decreasing trend with decreasing gravity.





P. Aeruginosa demonstrates significant growth under decreasing gravitational conditions. (Left Column) Centroid plot of cells within biofilm with larger circles indicating areas of greater cell density. (Middle Column) Overview of biofilm thickness density with warm colors indicating areas of greater thickness. (Right Column) Biofilm isosurface render demonstrating approximate substratum coverage and roughness.



- gravity was reduced.



I would like to thank the MSMHA program and BioServe Space Technologies for providing me the opportunity and resources necessary to pursue this project. Furthermore, I would like to personally thank Zeena Sadaf Nisar, University of Colorado Boulder candidate for B.S. 19', for helping me with this project. Finally, I would like to thank my peers and my committee members Dr. Salcedo and Dr. Finger for providing the moral support and advice necessary to get me to the finish line. This material is based upon work supported by the National Aeronautics and Space Administration under Grant No. 80NSSC17K0036



Donlan, Rodney M. "Biofilms: Microbial Life on Surfaces." Emerging Infectious Diseases, vol. 8, no. 9, 2002, pp. 881–890., doi:10.3201/eid0809.02006 2Hall-Stoodley, Luanne, and Paul Stoodley. "Evolving Concepts in Biofilm Infections." Cellular Microbiology, vol. 11, no. 7, 2009, pp. 1034–1043., doi:10.1111/j.1462-5822.2009.01323.x ³Khan, Hassan Ahmed, et al. "Nosocomial Infections and Their Control Strategies." Asian Pacific Journal of Tropical Biomedicine, vol. 5, no. 7, 2015, pp. 509–514 ⁴Kim, Dong-Ju, et al. "Relation of Microbial Biomass to Counting Units for Pseudomonas Aeruginosa." African Journal of Microbiology Research, vol. 6, no. 21, Sept. 2012, doi:10.5897/ajmr10.902 ⁵Kim, Wooseong, et al. "Spaceflight Promotes Biofilm Formation by Pseudomonas Aeruginosa." PLoS ONE, vol. 8, no. 4, 2013, doi:10.1371/journal.pone.0062437. [®]Vorregaard, M., Comstat2 - a modern 3D image analysis environment for biofilms, in Informatics and Mathematical Modelling. 2008, Technical University of Denmark

Complete list of references available in written manuscript

P. aeruginosa proliferated with greater capacity in both simulated Lunar and Martian environments, but the simulated microgravity environment was by far the worst with final cell concentrations increasing by over 72% compared to 1g controls.

Gravitational regime plays a role in the formation of *P. aeruginosa* biofilms and altering architectural characteristics. Most notably, the loss of microcolonies - indicated by decreasing roughness coefficients and increasing substratum coverage – was observed as simulated

Silicone loses its biofilm resistive capacities with decreasing simulated gravity.

Further studies are needed to elucidate the underlying mechanisms of these changes and determine if any of these changes may have detrimental effects to long-term space flight missions and eventual Lunar and Martian colonies.

ACKNOWLEDGEMENTS