

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/331499891>

# Patterns of *Clinostomum marginatum* infection in fishes and amphibians: Integration of field, genetic, and experimental approaches

Article in *Journal of Helminthology* · March 2019

DOI: 10.1017/S0022149X18001244

CITATIONS

2

READS

468

7 authors, including:



**Dana M Calhoun**

United States Geological Survey

30 PUBLICATIONS 142 CITATIONS

[SEE PROFILE](#)



**Katie Leslie**

University of Washington Seattle

3 PUBLICATIONS 4 CITATIONS

[SEE PROFILE](#)



**Tawni B Riepe**

Colorado State University

7 PUBLICATIONS 9 CITATIONS

[SEE PROFILE](#)



**Tyler Achatz**

University of North Dakota

10 PUBLICATIONS 15 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Black Spot Syndrome [View project](#)



Transmission of *Renibacterium salmoninarum* in Colorado native greenback cutthroat trout [View project](#)

## Research Paper

**Cite this article:** Calhoun DM, Leslie KL, Riepe TB, Achatz TJ, McDevitt-Galles T, Tkach VV, Johnson PTJ (2019). Patterns of *Clinostomum marginatum* infection in fishes and amphibians: integration of field, genetic, and experimental approaches. *Journal of Helminthology* 1–12. <https://doi.org/10.1017/S0022149X18001244>

Received: 15 November 2018

Accepted: 28 December 2018

**Key words:**

Amphibians; *Clinostomum* infection; fish; genetic analysis; host–parasite interactions; trematodes

**Author for correspondence:**

D.M. Calhoun

E-mail: [dana.calhoun@colorado.edu](mailto:dana.calhoun@colorado.edu)

# Patterns of *Clinostomum marginatum* infection in fishes and amphibians: integration of field, genetic, and experimental approaches

D.M. Calhoun<sup>1</sup>, K. L. Leslie<sup>1</sup>, T.B. Riepe<sup>1</sup>, T.J. Achatz<sup>2</sup>, T. McDevitt-Galles<sup>1</sup>, V.V. Tkach<sup>2</sup> and P.T.J. Johnson<sup>1</sup>

<sup>1</sup>Department of Ecology and Evolutionary Biology, University of Colorado, Ramaley N122 CB334, Boulder, CO 80309, USA and <sup>2</sup>Department of Biology, University of North Dakota, Grand Forks, ND 58202-9019, USA

**Abstract**

Digenetic trematodes of the genus *Clinostomum* are cosmopolitan parasites infecting fishes, amphibians, reptiles, and snails as intermediate hosts. Despite the broad geographical distribution of this genus, debate about the number of species and how they vary in host use has persisted. To better understand patterns of infection among host species and across life stages, we used large-scale field surveys and molecular tools to examine five species of amphibians and seven species of fishes from 125 California ponds. Among the 12,360 examined hosts, infection was rare, with an overall prevalence of 1.7% in amphibians and 9.2% in fishes. Molecular evidence indicated that both groups were infected with *Clinostomum marginatum*. Using generalized linear mixed effects models, host species identity and host life stage had a strong influence on infection status, such that *Lepomis cyanellus* (green sunfish) (49.3%) and *Taricha granulosa* (rough skinned newt) (9.2%) supported the highest overall prevalence values, whereas adult amphibians tended to have a higher prevalence of infection relative to juveniles (13.3% and 2.5%, respectively). Experimentally, we tested the susceptibility of two amphibian hosts (*Pseudacris regilla* [Pacific chorus frog] and *Anaxyrus boreas* [western toad]) to varying levels of cercariae exposure and measured metacercariae growth over time. *Pseudacris regilla* was 1.3× more susceptible to infection, while infection success increased with cercariae exposure dose for both species. On average, metacercariae size increased by 650% over 20 days. Our study highlights the importance of integrating field surveys, genetic tools, and experimental approaches to better understand the ecology of host–parasite interactions.

**Introduction**

*Clinostomum* (Digenea, Clinostomidae) is a cosmopolitan genus of digenetic trematodes prevalent in freshwater and estuarine aquatic environments worldwide. Its life cycle includes several stages and three different hosts (Osborn, 1911; Schell, 1985; McAllister *et al.*, 2010). Adult flukes live in the digestive tract, oesophagus, pharynx and/or mouth of fish-eating birds, typically Charadriiformes and Ciconiiformes (Kanev *et al.*, 2002), or, less commonly, within reptiles, mammals, or even humans (Yamashita, 1938; Belliappa, 1944; Witenberg, 1944; Ortlepp, 1963; Hara *et al.*, 2014). To date, various species of *Clinostomum* have been reported from over 40 species of birds from six continents, from great blue herons (*Ardea herodias herodias*) to African darters (*Anhinga rufa*) (supplementary table 1). Eggs are released by definitive hosts into an aquatic environment, where they hatch into miracidia and develop into sporocysts upon penetration into rams horn snails (typically of the genus *Helisoma*) (Cort, 1913; Hunter and Hunter, 1935; Smyth and Smyth, 1980). Members of other snail genera and families (e.g. *Lymnaea*, *Radix*, *Bulinus*, *Biomphalaria*) have also been reported as hosts, albeit less commonly (supplementary table 2; Esch *et al.*, 2001; Caffara *et al.*, 2011; Pinto *et al.*, 2015; Rosser *et al.*, 2017). Sporocysts release free-swimming cercariae that can infect a broad range of amphibian and freshwater fish species (supplementary table 3; Hoffman, 1999; McAllister *et al.*, 2010), within which they form loosely encysted metacercariae (Kanev *et al.*, 2002). Metacercariae can live in second intermediate hosts for up to four years (Elliott and Russert, 1949) and develop into adults once a suitable definitive host consumes the infected amphibian/fish host (Kalantan and Arfin, 1987; Bruni and Angelini, 2016).

Although trematodes in the genus *Clinostomum* are broadly distributed geographically and have been studied extensively, the taxonomy of this group has been the subject of extensive debate and discussion. Thus far, there are 31 recognized species of *Clinostomum* recorded from all continents other than Antarctica (see supplementary tables 1–3), of which 20 include some degree of support using genetic techniques (Locke *et al.*, 2015; Caffara *et al.*, 2017; Briosio-Aguilar *et al.*, 2018; Rosser *et al.*, 2018; Sereno-Urbe *et al.*, 2018). For the 14 species

reported in North America (see supplementary table 3), differentiation among species is often believed to differ based on their use of second intermediate hosts, with nine species infecting only fishes, one infecting only amphibians, and two using both host groups. Cort (1913) suggested that *Clinostomum* from frogs (*C. attenuatum*) differed sufficiently from those found in fishes (*C. marginatum*) to be considered a distinct species, which he hypothesized matured in different definitive hosts (bitterns and herons, respectively). This division was supported subsequently by several other researchers (Bear, 1933; Yamaguti, 1933; Hunter and Hunter, 1935; McAllister et al., 2010), although recent studies provide evidence that some *Clinostomum* species infect both fish and amphibians (McAllister, 1990; McAllister et al., 2010; Bonett et al., 2011; Caffara et al., 2011; Locke et al., 2015). Bonett and colleagues (2011) detected *C. marginatum* in both fishes (*Micropterus salmonoides*) and salamanders (*Eurycea tynerensis* and *E. spelaea*) in Oklahoma. While DNA sequence analyses have helped to validate the taxonomic identity, differentiated between species, and uniqueness of several previously identified species of *Clinostomum*, both in the New and Old World (Caffara et al., 2011; Sereno-Urbe et al., 2013, 2018; Locke et al., 2015; Pinto et al., 2015; Pérez Ponce de León et al., 2016; Rosser et al., 2018), the extent of their specificity to a second intermediate host remains conjectural.

Metacercariae of *Clinostomum* increase substantially in size within the second intermediate host (Osborn, 1911; Hopkins, 1933). Over 20 weeks in the laboratory, for instance, Osborn (1912) reported that *C. marginatum* metacercariae in a bass (*Micropterus* sp.) grew to 4001 µm in length, which was nearly the size of an adult fluke (4530 µm). Similarly, Hopkins (1933) found that the average length of metacercariae in field-collected pirate perch (*Aphredoderus sayanus*) increased from 200 µm in summer to > 2500 µm by winter (> 10-fold increase). Perhaps as a result of its notable growth capacity and large body size, *Clinostomum* infections are associated with host pathology. In fisheries, this parasite has gained notoriety as the causative agent of 'yellow grub disease' (Szalai and Dick, 1988; Hoffman, 1999; Overstreet and Curran, 2004; Bullard and Overstreet, 2008; Wang et al., 2017). Infected fish are often rejected for consumption because of the large and conspicuously yellow metacercariae, which can escape their cysts to become mobile while freshly captured fishes are displayed to consumers (Paperna, 1991). In amphibians, *Clinostomum* metacercariae have been linked to muscular dystrophy and scoliosis in heavily infected individuals (Belló et al., 2000; Perpiñán et al., 2010). For instance, histological examination of *Ambystoma tigrinum* (tiger salamander) revealed multifocal intramuscular, subcutaneous and coelomic trematodiasis, with associated necrosis and inflammation (Perpiñán et al., 2010). In a summary of 64 amphibian morbidity events in the USA during 1996–2001, Green et al. (2002) reported that *Clinostomum* sp. infections were the second most common cause of morbidity, with infected animals often having large dermal nodules on their skin. In contrast, Bruni and Angelini (2016) reported no short-term effects of *C. complanatum* exposure on the survival or dermal appearance of newts (*Triturus carnifex*), emphasizing the importance of further research to understand the patterns of infection among hosts and their influence on host fitness.

Despite the ubiquity of *Clinostomum* spp. infections in aquatic systems, large-scale field surveys of infection patterns in multiple amphibian and/or fish species are relatively rare. In one of the largest previous surveys of amphibian hosts, Gray and colleagues

(2007) examined 64,310 amphibians non-invasively for gross external signs of infection by *Clinostomum attenuatum*, focusing on the presence of nodules visible through the skin. This included three species (*Bufo cognatus*, *Ambystoma tigrinum mavortium* and *Spea multiplicata*) and two age classes (adult and juvenile) in the Southern Great Plains of the USA. They found extremely low prevalence in all species in both land-use areas (cropland and grasslands), with only 249 amphibians showing visible signs of infection (0.4%). Other field studies have reported a higher prevalence of *Clinostomum* spp.; for instance, Miller et al. (2004) reported that 14 of 21 (66.7%) amphibians were infected with *C. attenuatum*, including 7/11 *Spea multiplicata*, 2/2 *Bufo cognatus*, and 5/8 *Ambystoma tigrinum mavortium*, in wetlands in Texas. In fishes, Wang et al. (2017) reported a 9.8% prevalence of *Clinostomum complanatum* metacercariae in 1503 hosts from four families in Taiwan; the highest infection loads occurred in fishes of the Cyprinidae family. Overall, these surveys illustrate that patterns of infection vary among host species, at least in some cases, with higher prevalence in fishes relative to amphibians (Elliott and Russert, 1949; McAllister, 1990; Gray et al., 2007; Bonett et al., 2011; Caffara et al., 2014; Wang et al., 2017). However, few studies have directly compared infections across multiple species and different host groups (fish versus amphibians) in similar habitats.

In this study, we explored patterns of infection prevalence, intensity and host species use by *Clinostomum* sp. in 125 ponds within the Bay Area region of California, USA. Using large-scale field surveys over a five-year period, we tested the influence of host species identity and life stage on infection prevalence among five species of amphibians (n = 11,364 individuals) and seven species of fishes (n = 996). We used a combination of morphological and molecular methods to reveal the identity of parasites isolated from fishes and amphibians. To complement our field studies we used an experimental approach to evaluate the susceptibility and sensitivity of two amphibian species to varying numbers of *Clinostomum* sp. cercariae under controlled conditions. Over a period of 40 days, we estimated infection success, metacercarial growth, and host survival and growth. This work has implications for understanding the ecological and taxonomic patterns of *Clinostomum* infections and pathology in amphibian and fish hosts.

## Materials and methods

### Field surveys of infection among species, sites and years

Between May and August 2012–2017, we collected amphibians and fishes from temporary and permanent freshwater ponds in the Bay Area of California (Alameda, Contra Costa, and Santa Clara counties). Wetlands were selected to represent a range of sizes and hydroperiods, while containing freshwater gastropods as well as amphibian and/or fish hosts. The majority of the wetlands were built or modified to support cattle grazing (Garone, 2011), and are now managed as part of regional and county parks, research reserves, or as private land. In general, selected sites ranged in perimeter from < 50 m to > 500 m, with depths ranging from 0.5 to 5.0 m, with a surrounding habitat of oak woodland and annual grassland (McDevitt-Galles and Johnson, 2018).

With regards to amphibians, our focus was on late-stage larvae and recently metamorphosed individuals, although we did examine adults opportunistically. During transects along the wetland

edge, we used a combination of dip nets and hand-capture to collect ~10 individuals of each amphibian species per pond. The majority of the amphibian transects were conducted from late June to early August, with the exception of *Anaxyrus boreas* transects, which were conducted from early to mid-June due to the earlier emergence of the species. For bullfrogs (*Lithobates catesbeiana*), we captured metamorphs and adults at night with the aid of a headlamp. For fishes, we used habitat-stratified dip net sweeps (45.7 cm D-frame with 1.2 mm mesh), hauls with one of two seine nets (1.2 × 1.8 m or 4.5 × 1.8 m, each with 3 mm mesh), and rod and reel to collect ~10 adult individuals per taxon per visit. We ensured that our sampling methods adequately captured host and parasite richness using rarefaction curves and richness estimators (Johnson *et al.*, 2013). See Johnson *et al.* (2013) for additional details on sampling methods. Fishes were sampled in years 2013 through 2017, whereas amphibians were sampled across all years (2012–2017).

After collection, we euthanized hosts humanely using buffered MS-222 (dose = 1 g/500 ml of water), measured their snout–vent length (SVL; amphibians) or total length (fishes) using digital calipers, and examined all major organs and tissues (e.g. heart, liver, intestines, rectum, stomach, tongue, mandible, kidneys, gills, and fat bodies) for macroparasites under an Olympus SZX10 stereo dissecting microscope (Olympus Corporation, Tokyo, Japan). Parasites were heat-killed and preserved in 70% ethanol for morphological examination and in 95% molecular-grade ethanol for molecular analysis, with subsequent storage at –20°C. We counted and identified the parasites under 60–200× magnification using the keys of Lehmann (1954), Schell (1985), Sleigh (1991), Hoffman (1999), Gibson *et al.* (2002), Duszynski *et al.* (2007) and Anderson *et al.* (2009) as well as characteristic morphological features. Specifically, *Clinostomum* spp. have a stout body, oral sucker surrounded by a collar that is smaller in size when compared to the ventral sucker, and a bulbous oesophagus with little to no pharynx (Schell, 1985; Hoffman, 1999; Caffara *et al.*, 2011). After organ removal, we inspected the body cavity, muscles and host skin tissue layers for larval parasites, such as encysted trematode metacercariae (see Calhoun *et al.*, 2018).

To statistically explore patterns of *Clinostomum* sp. infection among naturally occurring hosts, we subset our dataset pond-by-year combinations in which at least one host was infected, regardless of the host species. This included information on 4745 individual amphibian hosts representing five species (*A. boreas*, *P. regilla*, *L. catesbeiana*, *Taricha granulosa*, *T. torosa*) and 513 fishes representing five species (*Gambusia affinis*, *Micropterus salmoides*, *Lepomis macrochirus*, *L. microlophus*, *L. cyanellus*). For analyses of infection prevalence, we considered the unit of analysis to be a host population, defined here as a unique combination of pond, sample year, and host species. To assess how infection prevalence varied with host species and life stage, we used generalized linear mixed models (GLMMs) with a binomial distribution and a logit link while accounting for the pond-by-year sampling event as a random intercept term to address the non-independence of hosts collected from the same pond-by-year combination (Bolker *et al.*, 2009). As the response variable, we used the *cbind* function in R (R Core Team, 2014) to combine vectors for the number of infected and uninfected hosts in a given population (Crawley, 2002). This has the advantage of inherently accounting for sample size, such that more heavily sampled populations are appropriately weighted, and avoiding the need to set arbitrary sample size thresholds. For all

analyses, quantitative fixed effects were scaled prior to inclusion by subtracting the mean and dividing by the standard deviation. Models were built using the *glmer* function in the *lme4* package (R Core Team, 2014; Bates *et al.*, 2015; Kuznetsova *et al.*, 2017).

For analyses of amphibian hosts specifically, we included fixed effects for species identity and life stage (both as factors), the latter of which was divided between juveniles (late-stage larvae and recent metamorphs) and adults. For amphibians, the analysis included 189 populations from 51 pond-year combinations. To compare patterns of infection between amphibians and fishes, we included 300 fish and amphibian populations from 65 pond-years. Using a similar modelling approach to that described above, we tested how host type (fish vs amphibian) influenced infection prevalence, with host species identity initially incorporated as a random intercept term. After detecting an overall difference between fishes and amphibians, we used species identity as a fixed effect, followed by Tukey pairwise comparisons using the *glht* function in the *multcomp* package as a post-hoc analysis of which species pairs differed (Hothorn *et al.*, 2013; Bretz *et al.*, 2016). Our overall goals were to investigate prevalence variation between fish and amphibians, as well as among individual amphibian and fish species.

### Genetic analysis

We obtained genomic DNA from 11 samples of *Clinostomum* sp. obtained from various fish and amphibian hosts from different ponds in the study area and metacercariae collected from experimental amphibians (table 1). DNA was isolated according to Tkach and Pawlowski (1999) or using a ZR Genomic DNA™ Tissue Micro Prep (Zymo Research, Irvine, California, USA) following the manufacturer's instructions. Following preliminary morphological identification using Schell (1985) and Jones *et al.* (2005), we used a single *Clinostomum* metacercaria (or a fragment in the case of larger individuals) for each DNA extraction. DNA fragments of approximately 1140 base pairs and spanning the 3' end of 18S nuclear rDNA gene, complete internal transcribed spacer region (ITS1 + 5.8S + ITS2) and a few nucleotides at the 5' end of the 28S gene as well as the partial mitochondrial *cox1* gene covering the traditional 'barcoding' region, were amplified by polymerase chain reaction (PCR) on a BioRad T-100 (Hercules, California, USA) thermal cycler. Forward primer ITSF (5'-CGCCGTCGCTACTACCGATTG-3'), and reverse primer digl2r (5'-CCGCTTAGTGATATGCTT-3') from Tkach and Snyder (2007) were used for rDNA amplifications; forward primers *Cox1-Schist5'* (5'-TCTTTTRGATCATAAGCG-3') from Lockyer *et al.* (2003) and reverse primer *cox650r* (5'-CCAAAAACCAAACATATGCTG-3') from Kudlai *et al.* (2015) were used for *cox1* amplifications. We ran PCR reactions using OneTaq Master Mix (New England Biolabs, Ipswich, Massachusetts, USA) following the manufacturer's instructions, with an annealing temperature of 53°C for rDNA amplifications and 43°C for *cox1* amplifications. PCR products were purified using ExoSAP-IT (Affymetrix, Santa Clara, California, USA), and sequencing reactions were prepared using the PCR primers mentioned above as well as the internal reverse primer *d58r* (5'-CACGAGCCGAGTGATCCACCGC-3') designed by V. Tkach. Purified PCR products were cycle-sequenced using a BigDye terminator v. 3.1 cycle sequencing kit (Applied Bio Systems, Foster City, California, USA). Sequencing reactions were alcohol-precipitated and run on an ABI Prism 3130™ automated capillary sequencer. We assembled contiguous sequences using

**Table 1.** Hosts, collection locations (all in California) and GenBank accession numbers of the sequenced *Clinostomum marginatum* metacercariae. Scientific and common names of hosts were confirmed using <https://amphibiaweb.org/> and <http://www.fishbase.org/search.php>.

| Host scientific name         | Host common name      | Collection location      | Accession numbers |          |
|------------------------------|-----------------------|--------------------------|-------------------|----------|
|                              |                       |                          | rDNA              | cox1     |
| <i>Micropterus salmoides</i> | largemouth black bass | 37.6394572, -121.9419164 | MK424220          | MK426657 |
| <i>Taricha granulosa</i>     | rough skinned newt    | 37.624493, -122.004644   | MK424221          | –        |
| <i>Taricha torosa</i>        | California newt       | 37.624493, -122.004644   | MK424222          | –        |
| <i>Pseudacris regilla</i>    | Pacific tree frog     | 37.363084, -121.731918   | MK424223          | MK426658 |
| <i>Pseudacris regilla</i>    | Pacific tree frog     | 37.9222114, -122.218118  | MK424224          | MK426659 |
| <i>Pseudacris regilla</i>    | Pacific tree frog     | 37.3015, -121.67259      | MK424225          | MK426660 |
| <i>Micropterus salmoides</i> | largemouth black bass | 37.6394572, -121.9419164 | MK424226          | MK426661 |
| <i>Taricha torosa</i>        | California newt       | 37.9456526, -122.140267  | MK424227          | –        |
| <i>Pseudacris regilla</i>    | Pacific tree frog     | 37.64527893, -121.918319 | MK424228          | –        |
| <i>Micropterus salmoides</i> | largemouth black bass | 37.4445, -121.71701      | MK424229          | MK426662 |
| <i>Anaxyrus boreas</i>       | western toad          | 37.363084, -121.731918   | MK424230          | MK426663 |

Sequencher™ (GeneCodes Corp., ver. 4.1.4), and submitted to GenBank (see table 1 for accession numbers). Obtained sequences (n = 13) were compared with all previously published sequences of *Clinostomum* spp. (from hosts collected worldwide) available in GenBank using the BLAST search and alignments in MEGA7 software (Kumar et al., 2016) with the goal of determining which species of *Clinostomum* were infecting our collected hosts. Specifically, we were interested in determining if the occurrence of *Clinostomum* sp. in our dataset supported current taxonomic and life history views that *C. marginatum* and *C. complanatum* can be found in fishes and amphibians whereas *C. attenuatum* infects amphibians only.

### Multiple-dose exposure experiment of amphibian larvae

To assess how infection success varied between amphibian species, over time and as a function of cercariae dosage, we experimentally exposed larvae of two amphibian species to one of four dosages of *Clinostomum* sp. cercariae. We collected multiple egg masses of *P. regilla* and *A. boreas* from local field sites, allowed them to hatch, and maintained them separately by species in 40 l containers until larvae reached Gosner (1960) stage 28. At that point, randomly selected larvae were transferred individually into containers filled with 400 ml of treated water (UV-sterilized, carbon-filtered, dechlorinated tapwater; hereafter referred to as treated water) and maintained at 18–20°C. Larvae were fed *ad libitum* a diet consisting of equal parts TetraMin (Tetra, Melle, Germany) fish food and ground *Spirulina* flakes. We changed the water and containers twice per week and maintained the light : dark cycle at 12 : 12 hours.

To obtain cercariae for experimental exposures, we placed field-collected snails (*Helisoma trivolvis*) previously identified as infected with *Clinostomum* into 50 ml centrifuge tubes with 40 ml of treated water; released cercariae were harvested within 1–2 hours of peak shedding, which typically occurred around 12.00 and 14.00 (approximately six hours after sunrise; Wang et al., 2017). We randomly assigned each individual amphibian larva to one of five treatments: 0 cercariae (control; *P. regilla* n = 20, *A. boreas* n = 20), 20 cercariae (*P. regilla* n = 23, *A. boreas*

n = 20), 40 cercariae (*P. regilla* n = 40, *A. boreas* n = 20), 100 cercariae (*P. regilla* n = 20, *A. boreas* n = 10), or 200 cercariae (*P. regilla* n = 5, *A. boreas* n = 5). Collected cercariae were pipetted into individual containers with amphibian larvae and allowed to infect them over a period of 24 hours (the maximum lifespan of most cercariae; Schell, 1985), after which we increased the water volume in the containers to 800 ml. Animals in the control treatment were ‘sham’ exposed to a similar amount of treated water without cercariae. At three time points (1.5 days [36 hours], 20 days and 40 days post exposure [*P. regilla* only]), we humanely euthanized a subset of animals, measured their SVL, and quantified the number of *Clinostomum* metacercariae using systematic necropsy. At each time point, a subset of metacercariae (n = 30) was photographed and measured to estimate parasite growth over time. Specifically, we heat-killed each metacercaria after mechanical excystment and measured it under a coverslip with a calibrated ocular micrometer on an Olympus B51 compound microscope (Olympus Corporation, Tokyo, Japan).

To evaluate the effects of *Clinostomum* cercariae exposure on host survival, we used a generalized linear model with a binomial distribution and logit link to test how host species (*P. regilla* vs *A. boreas*), exposure dosage (0, 20, 40, 100 or 200 cercariae), and their interaction influenced the likelihood a host survived to 20 days post exposure. We were not able to include hosts examined at 1.5 days post exposure because no infection was detected in any of those hosts (n = 3 *P. regilla* and 3 *A. boreas*). Among individuals that survived to 20 days, we used a general linear model (GLM) with a Gaussian distribution and an identity link to test for effects of host species, dosage, and their interaction on body size (SVL). To understand how parasite load (the number of detected metacercariae) varied by treatment, we used a GLMM with fixed effects for host species (as a factor), exposure dosage (as a numeric term), and their interaction (non-significant interactions were removed and models re-run). Because parasites per host were overdispersed, we used an overdispersed Poisson model by including an observation-level random effect (in this case the host identity), which functions similarly to a negative binomial model (Crofton, 1971; Lawless, 1987). Only exposed hosts that survived the 20-day trial were included. While we

**Table 2.** The overall total average infection prevalence, average intensity, and infection range ( $\pm 1$  SE) of *Clinostomum marginatum* detected in five amphibians and seven fish across the five-year study in freshwater ponds in California. Total number of hosts infected with *Clinostomum marginatum*.

| Species                         | Number dissected | Infection prevalence (%) | Average intensity ( $\pm 1$ SE) | Infection range |
|---------------------------------|------------------|--------------------------|---------------------------------|-----------------|
| <i>Anaxyrus boreas</i> *        | 1305             | 3.0                      | 0.24 $\pm$ 0.07                 | 1–44            |
| <i>Pseudacris regilla</i> *     | 4290             | 1.4                      | 3.63 $\pm$ 5.62                 | 1–30            |
| <i>Lithobates catesbeiana</i> * | 545              | 2.6                      | 3.38 $\pm$ 1.09                 | 2–7             |
| <i>Taricha granulosa</i> *      | 1398             | 3.3                      | 3.04 $\pm$ 0.36                 | 1–11            |
| <i>Taricha torosa</i> *         | 3826             | 1.1                      | 3.50 $\pm$ 0.60                 | 1–16            |
| <i>Carassius auratus</i>        | 41               | 0.0                      | –                               | –               |
| <i>Gambusia affinis</i> *       | 462              | 0.7                      | 1.43 $\pm$ 2.08                 | 1–3             |
| <i>Ictalurus</i> sp.            | 1                | 0.0                      | –                               | –               |
| <i>Lepomis cyanellus</i> *      | 53               | 1.9                      | 7.00 $\pm$ 1.53                 | 1–19            |
| <i>Lepomis macrochirus</i>      | 258              | 2.3                      | 2.22 $\pm$ 0.71                 | 1–7             |
| <i>Lepomis microlophus</i> *    | 7                | 0.1                      | 11 $\pm$ 0.00                   | 1–11            |
| <i>Micropterus salmoides</i>    | 174              | 4.2                      | 6.45 $\pm$ 0.91                 | 1–30            |

\* New host species record.

expected exposure dosage to positively affect the number of detected metacercariae, we sought to evaluate how infection success (number of metacercariae detected relative to number of cercariae administered) varied with exposure dosage, which can reveal evidence of density-dependent establishment. We therefore included an offset term for dosage, such that our analysis effectively tested how treatments affected the proportion of cercariae successful in establishing in each host (although this could also be done using the calculated proportional success of cercariae, the use of an offset term more effectively preserves the discrete nature of the response variable as a count of parasites).

Finally, we compared *P. regilla* hosts exposed to 40 cercariae and then examined at either 20 days post exposure ( $n = 19$ ) or 40 days post exposure ( $n = 14$ ) to test how time post exposure influenced the number of metacercariae detected. The modelling approach was similar to that described above (overdispersed Poisson GLMM), but the only fixed effect was days post exposure. Among a subset of *P. regilla* examined at either 20 or 40 days post exposure, we determined the effect of time (20 vs 40 days) and dosage (20, 40, 100, 200 cercariae) on metacercariae size using a linear model. As the response variable, we calculated metacercariae area using the formula for an ellipse and  $\log_{10}(x + 1)$ -transformed these values before analysis. Results were comparable using metacercariae length.


## Results

### Field surveys

Over six years (2012–2017) we examined 11,364 amphibians (1305 *A. boreas*, 4290 *P. regilla*, 545 *L. catesbeiana*, 1398 *T. granulosa*, and 3826 *T. torosa*) from 125 ponds, many of which were visited over multiple years. This collection also included 996 fishes (462 *G. affinis*, 174 *M. salmoides*, 258 *L. macrochirus*, 53 *L. cyanellus*, 41 *C. auratus*, 7 *L. microlophus*, and 1 *Ictalurus* sp.) from 25 ponds. Overall, *Clinostomum* infection prevalence was low but varied among host species, sites and sample year. For amphibians, 198 individuals were positive for *Clinostomum* (198/11,364; 1.7%), including 3.3% of *T. granulosa* (46/1398),

3.0% of *A. boreas* (39/1305), 2.6% of *L. catesbeiana* (14/545), 1.4% of *P. regilla* (58/4290) and 1.1% of *T. torosa* (41/3826). Of fishes, 9.2% were positive (92/997), including 24.1% of *M. salmoides* (42/174), 8.9% of *L. macrochirus* (23/258), 35.8% of *L. cyanellus* (19/53), 1.5% of *G. affinis* (7/462), and 14.3% of *L. microlophus* (1/7) (table 2). Neither *C. auratus* nor *Ictalurus* sp. supported *Clinostomum* infections, although these species had lower sample sizes ( $n = 41$  and  $n = 1$ , respectively, each from only a single study pond). Overall, *Clinostomum* intensity (only infected animals  $\pm 1$  SE) was low ( $4.1 \pm 0.3$  SE) and varied among host type, whereas fishes contained more metacercariae than amphibians across life stages (average intensity  $5.2 \pm 0.60$  SE;  $n = 92$ ; range 1–30 compared with  $3.5 \pm 0.3$  SE;  $n = 198$ ; 1–30, respectively; see table 2 for species-level average intensity and metacercariae range).

Among the 189 amphibian populations with at least one infected host (i.e. unique pond, sample year, host species combinations), both host species identity and life stage significantly influenced infection prevalence. Host species identity had a strong effect on infection (likelihood ratio test comparing models with and without host species, chi-square = 27.81,  $df = 4$ ,  $P < 0.0001$ ). Populations of *T. granulosa* supported the highest average prevalence (9.2%;  $n = 82$  pond-years; 565 individuals dissected), followed by *A. boreas* (5.1%;  $n = 63$  pond-years; 615 individuals dissected) (fig. 1). Although *T. torosa* and *P. regilla* were among the most commonly encountered amphibian species, they supported the lowest average prevalence values (2.9%;  $n = 180$  pond-by-year combinations; 1483 individuals dissected and 3.0%;  $n = 169$  pond-by-year combinations; 1793 individuals dissected, respectively) (fig. 1). Adult amphibians exhibited higher infection prevalence relative to juveniles of the same species (Binomial GLMM: life stage [adult] =  $2.486 \pm 0.352$ ,  $P < 0.00001$ ) (fig. 2). For instance, the average prevalence among adult *T. granulosa* was  $33.7\% \pm 0.05$  SE (21 pond-years; 135 dissected individuals) compared with  $0.8\% \pm 0.03$  SE in juveniles (61 pond-years; 430 dissected individuals). Focusing on juvenile amphibians specifically (omitting adults, for which it is difficult to know if they completed larval development at the site of collection), *A. boreas* had higher prevalence than each of the other four



**Fig. 1.** Average percent infection prevalence ( $\pm 1$  SE) of *Clinostomum marginatum* for five species of amphibians and fishes across life stage by pond-by-year combination where an infected host was present. Number of individuals dissected is reported. The pond-year combination for each host was as follows: *Anaxyrus boreas* (ANBO)  $n = 63$ ; *Pseudacris regilla* (PSRE)  $n = 169$ ; *Lithobates catesbeiana* (LICA)  $n = 37$ ; *Taricha granulosa* (TAGR)  $n = 82$ ; *T. torosa* (TATO)  $n = 180$ ; *Gambusia affinis* (GAAF)  $n = 21$ ; *Lepomis cyanellus* (LECY)  $n = 6$ ; *L. macrochirus* (LEMA)  $n = 24$ ; *L. microlophus* (LEMI)  $n = 2$ ; and *Micropterus salmoides* (MISA)  $n = 19$ . Both *Carassius auratus* and *Ictalurus* sp. supported no *C. marginatum* infection. Images from <http://aquaticparasites.org/>

species, whereas *P. regilla* had more infection relative to *T. torosa* (Tukey pairwise comparisons, all  $P < 0.05$ ). No other pairwise comparisons were significantly different ( $P > 0.05$ ).


*Clinostomum* infection was much more common in fishes compared with amphibians (Binomial GLMM: host type [fish] =  $1.530 \pm 0.702$ ,  $P = 0.029$ ). Of the five species with detected infections, *L. cyanellus* had highest average prevalence (49.3%;  $n = 6$  pond-by-year combinations; 35 dissected individuals), followed by *M. salmoides* (27.2%;  $n = 19$  pond-by-year combinations; 127 dissected individuals), and *L. microlophus* (25.0%; two pond-by-year combinations; three dissected individuals, although note the low sample size for this species). *Lepomis macrochirus* and *G. affinis* both supported low average prevalence, despite being two of the most commonly encountered fish species (19.4%;  $n = 24$  pond-by-year combinations; 142 dissected individuals and 3.3%;  $n = 21$  pond-by-year combinations; 205 dissected individuals, respectively) (fig. 1). Tukey pairwise comparisons with both fish and amphibian species indicated that sunfishes (genus *Lepomis*, especially *L. macrochirus* and *L. cyanellus*) and *M. salmoides* had higher infection prevalence than each of the amphibian species, as well as *G. affinis*.

Upon dissection of hosts, *Clinostomum* metacercariae were detected in a broad range of organs and tissues. In amphibians, metacercariae were large and loosely encysted in translucent cysts, often located between the epidermis and dermis (encompassing the stratum corneum and stratum germinativum but above the adipose tissue and lymph spaces; Duellman and Trueb, 1994) and frequently forming large, easily noticeable

nodules. In other cases, metacercariae were more deeply embedded in the muscle layer just underneath the subcutaneous tissue layer, and were less conspicuous. In amphibians, over half of animals infected contained metacercariae located between the epidermis and dermis near the mandible (52.6%). Other infection locations in amphibian hosts included the body cavity, oesophagus, gills (juvenile), kidneys, eyes, liver, intestinal mesentery, mouth, small intestine, tail (found in *Taricha* spp.), tongue and urinary bladder. The location of metacercariae also tended to vary by host species. In *T. torosa* and *T. granulosa*, for instance, most metacercariae were found around the mandible, whereas within *L. catesbeiana* the primary locations of infection were between the epidermis and dermis near the tail reabsorption site or hind limbs. Among fishes, *Clinostomum* occurred throughout the fins, body cavity, muscle tissue, between epidermal and dermal tissues, gills, liver, intestine, mouth and operculum of infected hosts. Muscle tissue was the most common location for metacercariae, accounting for 51.1% of the encountered parasites. Similar to amphibians, metacercariae in fishes were easily visible because they formed large nodules, which often appeared yellow in colour.

#### Genetic analysis

In total, we obtained DNA sequences from eight *Clinostomum* specimens from amphibians and three from fish (see table 1 for GenBank accession numbers), including specimens obtained in experiments using cercariae from naturally infected snails. As the ITS2 region did not sequence well in five of the specimens,



**Fig. 2.** Average percent infection prevalence ( $\pm 1$  SE) of *Clinostomum marginatum* for four species of amphibians and two life stages (juvenile and adult) by pond-by-year combination where an infected host was present. Number of individuals dissected is reported. The pond-year-combination for each host was as follows: adult *Pseudacris regilla* (PSRE)  $n=7$  and juvenile  $n=139$ ; adult *Lithobates catesbeiana* (LICA)  $n=9$  and juvenile  $n=20$ ; adult *Taricha granulosa* (TAGR)  $n=12$  and juvenile  $n=43$ ; adult *T. torosa* (TATO)  $n=11$  and juvenile  $n=108$ . No adult *Anaxyrus boreas* were dissected, and therefore they are not reported here. Images from <http://aquaticparasites.org/>

we compared the more variable complete ITS1 regions with flanking parts of 18S and 5.8S genes for all 11 sequenced samples. Based on the 733 bp long alignment all *Clinostomum* metacercariae obtained from both fishes and amphibians were identical in ITS1 region with the exception of a single base difference in one individual. Our ITS1 sequences were identical to previously published sequences of adult and larval specimens of *Clinostomum marginatum* (Rudolphi, 1819) from multiple host species and locations available in GenBank (Caffara *et al.*, 2011; Sereno-Uribe *et al.*, 2013; Pérez Ponce de León *et al.*, 2016; Rosser *et al.*, 2017). We were able to successfully amplify and sequence partial *cox1* gene from seven of our extracts from different hosts and locations (table 1). Upon trimming to the length of the shortest sequence, the *cox1* alignment was 574 bp long. Based on comparison with previously published sequences available in GenBank our sequences clearly belonged to *C. marginatum*, based on levels of intraspecific variability of 0–2.3%. These metacercariae were found in both amphibian and fish hosts.


### Experimental exposures

Of the amphibian larvae exposed to *Clinostomum* cercariae (not including controls), 65.5% of *A. boreas* ( $n=36$ ) and 87.5% of *P. regilla* ( $n=69$ ) became infected with one or more metacercariae. Among infected amphibians, the intensity  $\pm 1$  SE of encysted metacercariae varied across taxa, with a mean of  $17.6 \pm 2.8$  (1–117) in *P. regilla* and  $6.8 \pm 1.2$  (1–25) in *A. boreas*. *Clinostomum* metacercariae were detected between the epidermis and dermis and as deep as muscle tissues in both host species, similar to our observations in field-collected hosts. Neither exposure dosage nor host species (or their interaction) significantly influenced the

likelihood a tadpole survived to 20 days post exposure (binomial GLM, all  $P > 0.5$ ). Survival was generally high and similar between exposed and control hosts; for *A. boreas*, 80% of control hosts ( $n=16$ ) and 88.5% of parasite-exposed hosts ( $n=46$ ) survived to 20 days, whereas for *P. regilla*, 70% of controls ( $n=10$ ) and 80% of exposed hosts ( $n=59$ ) survived. Among surviving hosts, only host species identity affected SVL, with no added influence of cercariae exposure dosage (LM, *P. regilla* coefficient =  $-1.41 \pm 0.249$ ,  $P < 0.0001$ ; dose =  $0.003 \pm 0.0023$ ,  $P = 0.246$ ). On average,  $\pm 1$  SE, *A. boreas* larvae were  $11.50 \pm 0.17$  mm ( $n=62$ ), compared with  $10.13 \pm 0.19$  mm for *P. regilla* ( $n=69$ ), regardless of parasite exposure.

Exposure dosage and host species jointly influenced the number of detected metacercariae 20 days post exposure (Poisson GLMM: *P. regilla* =  $1.066 \pm 0.175$ ,  $P < 0.00001$ ; scale (dose) =  $0.347 \pm 0.079$ ,  $P < 0.00001$ ) (fig. 3). Because this model included an offset for exposure, these results indicate that the proportion of parasites successful in establishing within a host increased with both cercariae dosage and with host species (*P. regilla* > *A. boreas*). Time post-exposure also strongly affected the number of detected metacercariae. Among a subset of hosts exposed to 20 cercariae and examined at 36 hours post exposure (three *P. regilla* and three *A. boreas*), no metacercariae were detected. Similarly, for hosts that died within 10 days of exposure (seven *P. regilla* and eight *A. boreas*), only a single *P. regilla* exposed to 200 cercariae showed evident signs of infection (with three metacercariae). Based on the analysis of *P. regilla* hosts exposed to 40 cercariae and examined at 20 or 40 days post exposure ( $n=25$  and 10, respectively), time had a strongly positive effect on parasite load (Poisson GLMM: scale (days) =  $0.449 \pm 0.128$ ,  $P = 0.0004$ ). More specifically, the average number of metacercariae per host





**Fig. 3.** The infection load ( $\pm 1$  SE) of *Clinostomum marginatum* metacercariae recovered at 20 days per infected species of amphibian, *Pseudacris regilla* (PSRE) and *Anaxyrus boreas* (ANBO), per dose of cercariae (20, 40, 100, 200). Number of individuals dissected is reported.

$\pm 1$  SE more than doubled, from  $6.21 \pm 1.24$  on day 20 to  $14.2 \pm 1.96$  on day 40, despite an identical initial exposure level. These patterns probably reflect increases in our ability to detect metacercariae with time, rather than a real increase in infection (given that all exposures occurred within the first 24 hours).

The average size of *Clinostomum* metacercariae increased between days 20 and 40 post exposure. The estimated area of metacercariae increased 650% during this period, from  $0.0125 \pm 0.0008 \text{ mm}^2$  to  $0.0812 \pm 0.0009 \text{ mm}^2$ , with no effect of exposure dosage (LM on  $\log_{10}(\text{area} + 1)$ :  $\text{scale}(\text{days}) = 0.400 \pm 0.033$ ,  $P < 0.00001$ ;  $\text{scale}(\text{dose}) = 0.0038 \pm 0.033$ ,  $P = 0.91$ ). The average length of measured metacercariae across doses  $\pm 1$  SE at 20 days was  $179.44 \pm 9.41 \mu\text{m}$ , with an average width of  $89.44 \pm 4.67$  ( $n = 9$ ). At 40 days ( $n = 13$ ), the average length and width were  $383.08 \pm 21.07 \mu\text{m}$  and  $263.46 \pm 17.81 \mu\text{m}$ , respectively.

## Discussion

### Patterns of infection among species and over time in California

We detected *Clinostomum marginatum* infections in 10 host species (five amphibian and five fish species) over a five-year period of field sampling in the Bay Area of California, of which eight represent new host records (table 2). Our field surveys highlighted the importance of both host species and life stage in influencing parasite occurrence and prevalence. Overall, infection prevalence was low, with only 1.7% of amphibian and 9.2% of fish hosts showing infection. Of the 125 ponds sampled, many of which were visited in multiple years (average of three visits), only 40 supported *C. marginatum* in one or more host species. Fish hosts tended to exhibit a higher prevalence and intensity relative to amphibians, even at sites where both groups co-occurred (1.6 $\times$  higher). Among the amphibians, a somewhat surprising result was that adults tended to support a higher likelihood of *Clinostomum* infection relative to larval amphibians, particularly for the two species of *Taricha* (fig. 2) (average prevalence in adults = 22.2% and in juveniles = 1.7%). For waterborne parasites such as *Clinostomum* cercariae, we expected that most amphibians post metamorphosis would experience lower infection risk

(presence or load), given that contact with water is maximized during larval development, although there is still potential for infection risk. Even though we cannot be certain what drives the elevated level of *C. marginatum* infections in adult amphibians, this result could be a function of detection (as trematode metacercariae grow with age they may be easier to detect; see experimental results) or changes in properties of the adult amphibian skin, which has been demonstrated in experimental studies with other trematodes to be less susceptible to cercariae invasion (Johnson et al., 2011).

The patterns of infection observed in the current study are broadly similar to previous surveys for both amphibians and fishes. Fish supporting a higher prevalence of *C. marginatum* is consistent with the longer lifespan and typically larger body size of fishes (Poulin and Mouritsen, 2003; Calhoun et al., 2018). For instance, channel catfish can live for as long as 15 years, often leading to high levels of 'yellow grub' infection and metacercariae persisting for long periods (see Edney, 1940). Based on our field results, *T. granulosa* supported the highest average parasite prevalence (9.2%), followed by *A. boreas* (5.1%). In parallel, Gray et al. (2007) detected more infection by *Clinostomum attenuatum* in salamanders (*Ambystoma tigrinum mavortium*, 0.5%) compared with toads (*Bufo cognatus*, 0.4%) or frogs (*Spea multiplicata*, 0.4%). However, the *Clinostomum* detection method used by Gray et al. (2007) differed from our study, as they only inspected the epidermis of the animals externally for parasites, which could explain the higher prevalences in our study (we often detected parasites deep in the muscle tissues). In a field study from Illinois, Crawford and Kuhns (2008) found substantially higher *Clinostomum* spp. prevalence among two larval salamander species (33.3% in *A. jeffersonianum* and 6.8% in *A. texanum*), and Kuperman et al. (2004) reported that 70% of non-native African clawed frogs (*Xenopus laevis*) were infected in southern California, with a range of 1–4 metacercariae ( $n = 230$  hosts). With respect to fish hosts, we found an overall infection prevalence of 9.2% ( $n = 996$ ), which aligns closely with the 9.8% prevalence reported by Wang et al. (2017) in a study of river fishes in Taiwan. They found Cyprinidae fishes supported the highest average intensity; *Candidia barbata* supported the

highest average intensity (14.4; 1–107;  $n = 31$ ), in parallel to our findings for *L. cyanellus* ( $7.0 \pm 1.5$  SE; 1–19;  $n = 18$ ). Also, similar to our findings, previous work has detected a low level of macro-parasite taxonomic richness and load within the invasive mosquitofish (3.3% prevalence in the current survey). Over four years, Calhoun *et al.* (2018) reported that *G. affinis* supported 42–78% fewer parasite taxonomic groups and 97% fewer total parasites compared with other fishes in California ponds. In a study of *Clinostomum marginatum* in fishes from eastern Kansas, Klaas (1963) found bullhead catfishes to be the most commonly infected fishes, followed by *M. salmoides* (7.2%), *L. macrochirus* (4.4%) and *L. cyanellus* (1.7%) (see also Becker and Cloutman, 1975). We did not find infection in either *C. auratus* or *Ictalurus* sp., which have been reported to support infections. This is probably because both fishes were collected from a single pond over several years. Thus, if that site did not support infection in frogs or newts, we would not be able to assess for these species.

Similar to our study, Bonnett and colleagues (2011) analysed the prevalence, intensity and genetic diversity of *Clinostomum* spp. in fishes (*M. salmoides*;  $n = 34$ ) and salamanders (*Eurycea tynerensis*;  $n = 74$ ) in an Oklahoma pond and adjacent stream. They detected *C. marginatum* in both species, complementing the current findings from California. Moreover, they reported a higher infection prevalence and intensity in fish hosts relative to amphibians (91% of *M. salmoides* and 56% of *E. tynerensis*). Largemouth black bass exhibited a mean intensity of  $5.0 \pm 4.7$  SD metacercariae, compared with  $2.2 \pm 1.9$  SD for infected salamanders.

### Genetic insights into parasite taxonomy

Based on both nuclear rDNA and mitochondrial *cox1* sequences the samples obtained in this study belonged to *C. marginatum*. We detected *C. marginatum* infections in both amphibians and fishes from multiple freshwater ponds in California, including samples obtained using naturally infected snails from those ponds. We point out, however, that the total number of sequenced samples in this study was low, and future efforts to expand on this foundation will be essential to evaluate whether additional species co-occur. Previous molecular analyses have provided some resolution regarding the taxonomic status of species within the genus *Clinostomum* (Dzikowski *et al.*, 2004; Caffara *et al.*, 2011; Locke *et al.*, 2015; Pérez Ponce de León *et al.*, 2016; Sereno-Urbe *et al.*, 2018). Using 18S rRNA isolates of *C. complanatum* from Israel and *C. marginatum* from the USA, Dzikowski and colleagues (2004) demonstrated the validity of the two species as they differed at 22 bases. More recently, Caffara *et al.* (2011) used the genetic markers ITS1 and COI, which have been determined to be more specific genes to differentiate species (compared with 18s rRNA alone), as well as morphological characteristics of metacercariae and adults to argue that *C. marginatum* is a separate species from *C. complanatum*. These findings support the contention of a geographical distinction between the American and European species. Using similar methods, Sereno-Urbe *et al.* (2013) used both mitochondrial and ribosomal markers on specimens from freshwater fishes and birds in Mexico to validate *C. complanatum*, while also identifying a new species: *Clinostomum tataxumui*. A large-scale survey of *Clinostomum* specimens from amphibian and fish hosts across New World and Old World regions further supported the geographical separation of species (Locke *et al.*, 2015), although without a clear separation between infections of amphibian and fish

hosts. Pinto *et al.* (2015) argued that the diversity of *Clinostomum* species is probably underestimated, based on a molecular study of cercariae from *Biomphalaria* spp. in Brazil. This opinion is supported by the most recent descriptions of four additional new species from North America (Briosio-Aguilar *et al.*, 2018; Rosser *et al.*, 2018).

Despite recent explorations of *Clinostomum* genetic history, few studies have examined species-level identifications within the same system in comparison of intermediate fish and amphibian hosts. Our study emphasized the importance of genetic data to identify species of *Clinostomum* in aquatic systems where multiple species of intermediate hosts frequently co-occur. Although classical studies contend that species identity varies as a function of intermediate host identity (amphibians vs fishes), our findings indicated that *C. marginatum* often infects amphibians as well as fishes in the same habitats. Prior to recent molecular studies (Caffara *et al.*, 2011; Sereno-Urbe *et al.*, 2013, 2018; Pérez Ponce de León *et al.*, 2016; Rosser *et al.*, 2016, 2017), *Clinostomum* species determination was often based on geographical location (New vs Old World) and intermediate host identity (fishes vs amphibians) (Yamaguti, 1933; Hunter and Hunter, 1935; McAllister, 1990). Alongside the findings of Bonnett *et al.* (2011), our results indicate that *C. marginatum* frequently infects both amphibians and fishes that co-occur in the same aquatic environment.

Although there have been multiple reports of *C. marginatum* infections in fish intermediate hosts, our results highlight three new host records in fishes (*G. affinis*, *L. cyanellus*, *L. microlophus*) and five new host records in amphibians (*A. boreas*, *P. regilla*, *L. catesbeiana*, *T. granulosa*, *T. torosa*) (table 2). Most of the amphibian host species in our study have previously reported infections by either *Clinostomum attenuatum* or *Clinostomum* spp. (supplementary table 3). For instance, *P. regilla* and *L. catesbeiana* have previously supported infection by *Clinostomum* spp. (Muzzall, 1991; Lemke *et al.*, 2008) and *C. attenuatum* (Cort, 1913; Jinks and Johnson, 1970). These results collectively indicate that the use of second intermediate host identity alone is probably insufficient to differentiate the species of *Clinostomum*, and therefore the previously suggested high degree of intermediate host specificity should be interpreted with caution.

### Experimental results for parasite growth and host pathology

Based on our experimental results, metacercariae of *C. marginatum* grew approximately 6.5-fold within *P. regilla* host over 20 days. This finding is notable in that metacercariae of most trematode species tend to grow relatively little within second intermediate hosts (Schell, 1985). For instance, *Fibricola seolensis* recovered from experimentally infected tadpoles grew 116% over 20 days (Mitchell *et al.*, 2002), whereas *Centrocestus formosanus* from infected fishes grew by 54% over 50 days. Others have even reported a decrease in growth; the width of *Ornithodiplostomum ptychocheilus* metacercariae in experimentally infected *Pimephales promelas* (fathead minnows) decreased by 31% (Sandland and Goater, 2000). Species of *Clinostomum* exhibit remarkably high levels of growth within intermediate hosts, in part associated with characteristics of the parasites' tegument (Lo *et al.*, 1985; Abidi *et al.*, 1988; Aho, 1990). For instance, Lo *et al.* (1985) noted that *C. complanatum* metacercariae had blood vessels penetrating the cysts, whereas in *C. complanatum* Abidi *et al.* (1988) reported a stretching capability in the tegument of metacercariae. Both of these studies suggest that increased access to host nutrients

probably assists in the growth and development of *Clinostomum marginatum*.

Despite this remarkable growth in metacercariae, we found no evidence of pathological effects in amphibian larvae exposed to *Clinostomum* cercariae, even at relatively high levels of exposure (200 cercariae). There was no difference in survival between exposed and unexposed hosts, and even among infected individuals there was no obvious effect on host growth. This result is consistent with previous studies showing little damage of initial exposure dosage on host fitness (Klaas, 1963), although other studies have reported pathology such as localized haemorrhages associated with infection (Elliott and Russert, 1949; Adeyemo and Agbede, 2008; Roberts, 2012). For instance, *Clinostomum tilapiae* in infected fishes causes inflammation and lesions, in some cases leading to host death (Adeyemo and Agbede, 2008). Similarly, Roberts (2012) found that fishes infected with clinostomatids had inhibited growth and increased weight loss. Given that we examined hosts over a limited time period (20 days), it is possible that more significant pathology would have occurred over longer time periods (including early larval development and/or especially the transition to metamorphosis).

Another interesting finding of the experimental study was that hosts exhibited density-dependent infection success, such that infection success (the percentage of administered cercariae that formed metacercariae) increased monotonically with exposure dosage. For example, at a dosage of 40 cercariae, *P. regilla* supported an average  $\pm 1$  SE of  $8.7 \pm 1.2$  metacercariae per host (21.7% infection success), whereas infection success increased to 30.7% when hosts were exposed to 100 cercariae (metacercariae count of  $30.7 \pm 5.1$ ). In contrast, Johnson *et al.* (2012) exposed 12 species of amphibians to a range of *Ribeiroia ondatrae* cercariae (0–200) and reported that the proportion of established metacercariae (relative to the number of cercariae administered) was independent of exposure dose. Similar results are consistently found in the *Microphallus* spp. trematode system in multiple dose exposures in snails (*Potamopyrgus antipodarum*) (Osnas and Lively, 2004) and in cercariae to amphipods (*Gammarus insensibilis*) (Brown *et al.*, 2003). Whereas many studies have reported that infection success decreases with exposure dosage (Koprivnikar *et al.*, 2017; Orlofske *et al.*, 2017), the ecological or physiological mechanisms underlying an increase in parasite establishment with density remain conjectural. The explanation for this trend could be twofold. In part, results could reflect an increase in the detection of metacercariae over time. For the subset of animals dissected at 36 hours post infection, no metacercariae were detected; however, at 20 days post infection, *P. regilla* hosts supported an average load of 9.5 metacercariae across treatments, highlighting probable differences in researchers' ability to observe metacercariae (as a function of growth within the host). Secondly, parasite performance could also be dose dependent, such that hosts exposed to higher levels of cercariae could exhibit a reduced capacity to resist or clear infections by *Clinostomum*. Future studies that incorporate immune parameter measurements could effectively disentangle the biological mechanisms underlying these observations.

**Supplementary material.** To view supplementary material for this article, please visit <https://doi.org/10.1017/S0022149X18001244>

**Author ORCIDs.**  D.M. Calhoun, 0000-0002-9483-2064.

**Acknowledgements.** We gratefully acknowledge the following members of the field and dissection teams who assisted in collection and dissection of amphibians, fishes and snails: Wynne Moss, Katie Richgels, Jeremy

Henderson, Evan Esfahani, Mary Jade Farruggia, Dina Soltow, Aaron Klingborg, Megan Housman, Margaret Summerside, Neal Handloser, Dane McKittrick, Sarah Goodnight, Sara Paull, Jay Bowerman, Sarah Kupferberg, Emily Hannon, Christina Garcia, Jackie Gregory, Bryan LaFonte, Keegan McCaffrey and Audrey Omeimrin. We thank two reviewers and the editor for their comments on the final manuscript.

**Financial support.** This work was supported through funding from the National Institutes of Health (R01-GM109499), National Science Foundation (DEB-1149308), and the David and Lucile Packard Foundation.

**Conflict of interest.** None.

**Ethical standards.** The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national and institutional guides on the care and use of laboratory animals.

## References

- Abidi SMA *et al.* (1988) *Clinostomum complanatum*: tegumental surface changes during in vivo development. *International Journal for Parasitology* **18**, 433–439.
- Adeyemo AO and Agbede SA (2008) Histopathology of tilapia tissues harbouring *Clinostomum tilapiae* parasites. *African Journal of Biomedical Research* **11**, 115–118.
- Aho JM (1990) Helminth communities of amphibians and reptiles: comparative approaches to understanding patterns and processes. In Esch GW, Bush AO and Aho JM (eds), *Parasite Communities: Patterns and Processes*. London: Chapman and Hall, pp. 157–195.
- Anderson RC, Chabaud AG and Willmott S (2009) *Keys to the Nematode Parasites of Vertebrates: Archival Volume*. London: CABI.
- Bates D *et al.* (2015) Parsimonious mixed models. *arXiv preprint arXiv:1506.04967*.
- Bear J (1933) Note sur un nouveau trematode, *Clinostomum lophophallum* sp. nov. *Revue Suisse de Zoologie* **40**, 312–347.
- Becker DA and Cloutman DG (1975) Parasites of selected gamed fishes of Lake Fort Smith, Arkansas. *Journal of the Arkansas Academy of Science* **29**, 12–18.
- Belliappa AB (1944) On a species of *Clinostomum* in a cat. *Indian Veterinary Journal* **21**, 101–102.
- Belló ARR *et al.* (2000) Lipid peroxidation induced by *Clinostomum detrunctum* in muscle of the freshwater fish *Rhamdia quelen*. *Diseases of Aquatic Organisms* **42**, 233–236.
- Bolker BM *et al.* (2009) Generalized linear mixed models: a practical guide for ecology and evolution. *Trends in Ecology & Evolution* **24**, 127–135.
- Bonett RM *et al.* (2011) Distribution, abundance, and genetic diversity of *Clinostomum* spp. metacercariae (Trematoda: Digenea) in a modified Ozark stream system. *Journal of Parasitology* **97**, 177–184.
- Bretz F, Westfall P and Hothorn T (2016) *Multiple Comparisons Using R*. Boca Raton, FL: CRC Press.
- Brown SP *et al.* (2003) Field evidence for density-dependent effects in the trematode *Microphallus papillorobustus* in its manipulated host, *Gammarus insensibilis*. *Journal of Parasitology* **89**, 668–672.
- Briosio-Aguilar R *et al.* (2018) Morphological and molecular characterization of an enigmatic clinostomid trematode (Digenea: Clinostomidae) parasitic as metacercariae in the body of freshwater fishes (Cichlidae) across Middle America. *Journal of Helminthology*, <https://doi.org/10.1017/S0022149X18000445>.
- Bruni G and Angelini C (2016) No short term effect of *Clinostomum complanatum* (Trematoda: Digenea: Clinostomatidae) on survival of *Triturus carnifex* (Amphibia: Urodela: Salamandridae). *Acta Herpetologica* **11**, 75–79.
- Bullard SA and Overstreet RM (2008) Digeneans as enemies of fishes. *Fish Diseases* **2**, 817–976.
- Caffara M *et al.* (2011) Morphological and molecular differentiation of *Clinostomum complanatum* and *Clinostomum marginatum* (Digenea: Clinostomidae) metacercariae and adults. *Journal of Parasitology* **97**, 884–891.

- Caffara M et al.** (2014) Metacercariae of *Clinostomum complanatum* (Trematoda: Digenea) in European newts *Triturus carnifex* and *Lissotriton vulgaris* (Caudata: Salamandridae). *Journal of Helminthology* **88**, 278–285.
- Caffara M et al.** (2017) A morphological and molecular study of Clinostomid metacercariae from African fish with redescription of *Clinostomum tilapiae*. *Parasitology* **144**, 1519–1529.
- Calhoun DM, McDevitt-Galles T and Johnson PTJ** (2018) Parasites of invasive freshwater fishes and the factors affecting their richness. *Freshwater Science* **37**, 134–146.
- Cort WW** (1913) Notes on the trematode genus *Clinostomum*. *Transactions of the American Microscopical Society* **32**, 169–182.
- Crawford JA and Kuhns AR** (2008) Identifying the distribution and habitat of Jefferson's salamander, *Ambystoma jeffersonianum*, in Illinois. *Illinois Natural History Survey* **24**, 1–20.
- Crawley MJ** (2002) *Statistical Computing*. London: John Wiley & Sons.
- Crofton HD** (1971) A model of host–parasite relationships. *Parasitology* **63**, 343–364.
- Duellman WE and Trueb L** (1994) *Biology of Amphibians*. New York, NY: McGraw-Hill.
- Duszynski DW, Bolek MG and Upton SJ** (2007) Coccidia (Apicomplexa: Eimeriidae) of the amphibians of the world. *Zootaxa* **1667**, 1–77.
- Dzikowski R et al.** (2004) *Clinostomum complanatum* and *Clinostomum marginatum* (Rudolphi, 1819) (Digenea: Clinostomidae) are separate species based on differences in ribosomal DNA. *Journal of Parasitology* **90**, 413–414.
- Edney JM** (1940) The longevity of *Clinostomum marginatum* metacercariae in channel catfish. *Journal of the Tennessee Academy of Science* **15**, 380–388.
- Elliott AM and Russert LR** (1949) Some condition characteristics of a yellow perch population heavily parasitized by *Clinostomum marginatum*. *Journal of Parasitology* **35**, 183–190.
- Esch GW, Curtis LA and Barger MA** (2001) A perspective on the ecology of trematode communities in snails. *Parasitology* **123**, 57–75.
- Garone P** (2011) *The Fall and Rise of the Wetlands of California's Great Central Valley*. Berkeley, CA: University of California Press.
- Gibson DI, Jones A and Bray RA** (2002) *Keys to the Trematoda*. London: CAB International and Natural History Museum.
- Gosner KL** (1960) A simplified table for staging anuran embryos and larvae with notes on identification. *Herpetologica* **16**, 183–190.
- Gray MJ et al.** (2007) Influences of agricultural land use on *Clinostomum attenuatum* metacercariae prevalence in southern Great Plains amphibians. *Herpetological Conservation and Biology* **2**, 23–28.
- Green DE et al.** (2002) Epizootiology of sixty-four amphibian morbidity and mortality events in the USA, 1996–2001. *Annals of the New York Academy of Sciences* **969**, 323–339.
- Hara H et al.** (2014) Human laryngitis caused by *Clinostomum complanatum*. *Nagoya Journal of Medical Science* **76**, 181–185.
- Hoffman GN** (1999) *Parasites of North American Freshwater Fishes*. New York, NY: Cornell University Press.
- Hopkins SH** (1933) Note on the life history of *Clinostomum marginatum* (Trematoda). *Transactions of the American Microscopical Society* **52**, 147–149.
- Hothorn T et al.** (2013) Multcomp: simultaneous inference in general parametric models. R package version 1.2-18. R Foundation for Statistical Computing, Vienna, Austria.
- Hunter WS and Hunter GW** (1935) Studies on *Clinostomum*. II. The miracidium of *C. marginatum* (Rud.). *Journal of Parasitology* **21**, 186–189.
- Jinks JL and Johnson JC** (1970) Trematodes of *Rana catesbeiana* from three strip-mine lakes in southeast Kansas. *Transactions of the Kansas Academy of Science* **73**, 519–520.
- Johnson PTJ, Kellermanns E and Bowerman J** (2011) Critical window of disease risk: amphibian pathology driven by developmental changes in host resistance and tolerance. *Functional Ecology* **25**, 726–734.
- Johnson PTJ et al.** (2012) Living fast and dying of infection: host life history drives interspecific variation in infection and disease risk. *Ecology Letters* **15**, 235–242.
- Johnson PTJ et al.** (2013) Biodiversity decreases disease through predictable changes in host community competence. *Nature* **494**, 230–233.
- Jones A, Bray RA and Gibson DI** (2005) *Keys to the Trematoda*. London: CAB International and Natural History Museum.
- Kalantan A and Arfin M** (1987) Seasonal incidence and pathogenicity of the metacercariae of *Clinostomum complanatum* in *Aphanius dispar*. *Japan Journal of Parasitology* **36**, 17–23.
- Kanev I, Radev V and Fried B** (2002) Family Clinostomidae Lühe, 1901. In Gibson GI, Jones A and Bray RA (eds), *Keys to the Trematoda*, Vol 1. New York, NY: CAB Publishing, pp. 113–120.
- Klaas EE** (1963) Ecology of the trematode *Clinostomum marginatum* and its hosts in eastern Kansas. *Transactions of the Kansas Academy of Science* **66**, 519–538.
- Koprivnikar J et al.** (2017) Does spatial heterogeneity in host distribution drive parasite aggregation among hosts? *Oikos* **127**, 99–110.
- Kudlai O et al.** (2015) A new species of *Drepanocephalus* Dietz, 1909 (Digenea: Echinostomatidae) from the double-crested cormorant *Phalacrocorax auritus* (Lesson) (Aves: Phalacrocoracidae) in North America. *Systematic Parasitology* **90**, 221–230.
- Kumar S, Stecher G and Tamura K** (2016) MEGA7: molecular evolutionary genetics analysis version 7.0 for bigger datasets. *Molecular Biology and Evolution* **33**, 1870–1874.
- Kuperman BI et al.** (2004) Parasites of the African clawed frog, *Xenopus laevis*, in Southern California, USA. *Comparative Parasitology* **71**, 229–232.
- Kuznetsova A, Brockhoff PB and Christensen RHB** (2017) lmerTest package: tests in linear mixed effects models. *Journal of Statistical Software* **82**, 1–27.
- Lawless JF** (1987) Negative binomial and mixed Poisson regression. *Canadian Journal of Statistics* **15**, 209–225.
- Lehmann DL** (1954) Some helminths of west coast urodeles. *Journal of Parasitology* **40**, 231–238.
- Lemke LB et al.** (2008) Infestation of wild-caught American bullfrogs (*Rana catesbeiana*) by multiple species of metazoan parasites. *Journal of the American Association for Laboratory Animal Science* **47**, 42–46.
- Lo CF, Chen SC and Wang CH** (1985) The study of *Clinostomum complanatum* (Rud., 1814) V. The influences of metacercaria of *Clinostomum complanatum* on fish. *Fish Pathology* **20**, 305–312.
- Locke SA et al.** (2015) A large-scale molecular survey of *Clinostomum* (Digenea, Clinostomidae). *Zoologica Scripta* **44**, 203–217.
- Lockyer AE et al.** (2003) The phylogeny of the Schistosomatidae based on three genes, with emphasis on the interrelationships of *Schistosoma* Weinland, 1858. *Parasitology* **126**, 203–224.
- McAllister CT** (1990) Metacercaria of *Clinostomum complanatum* (Rudolphi, 1814) (Trematoda: Digenea) in a Texas salamander, *Eurycea neotenes* (Amphibia: Caudata), with comments on *C. marginatum* (Rudolphi, 1819). *Journal of the Helminthological Society of Washington* **57**, 69–71.
- McAllister CT et al.** (2010) Metacercariae of *Clinostomum* (Trematoda: Digenea) from three species of *Ambystoma* (Caudata: Ambystomatidae) from Arkansas and Illinois, USA. *Comparative Parasitology* **77**, 25–30.
- McDevitt-Galles T and Johnson PTJ** (2018) Drought attenuates the impact of fish on aquatic macroinvertebrate richness and community composition. *Freshwater Biology* **63**, 1457–1468.
- Miller DL et al.** (2004) Metacercariae of *Clinostomum attenuatum* in *Ambystoma tigrinum mavortinum*, *Bufo cognatus* and *Spea multiplicata* from west Texas. *Journal of Helminthology* **78**, 373–376.
- Mitchell CE, Tilman D and Groth JV** (2002) Effects of grassland plant species diversity, abundance, and composition on foliar fungal disease. *Ecology* **83**, 1713–1726.
- Muzzall PM** (1991) Helminth infracommunities of the frogs *Rana catesbeiana* and *Rana clamitans* from Turkey Marsh, Michigan. *Journal of Parasitology* **77**, 366–371.
- Orlowski SA et al.** (2017) Experimental investigation of alternative transmission functions: quantitative evidence for the importance of non-linear transmission dynamics in host–parasite systems. *Journal of Animal Ecology* **87**, 703–715.
- Ortlepp RJ** (1963) Clinostomid trematodes as aberrant parasites in the mouth of the domestic cat (*Felis catus domesticus*). *Onderstepoort Journal of Veterinary Research* **30**, 137–143.

- Osborn HL** (1911) On the distribution and mode of occurrence in the United States and Canada of *Clinostomum marginatum*, a trematode parasitic in fish, frogs and birds. *The Biological Bulletin* **20**, 350–366.
- Osborn HL** (1912) On the structure of *Clinostomum marginatum*, a trematode parasite of the frog, bass and heron. *Journal of Morphology* **23**, 189–229.
- Osnas EE and Lively CM** (2004) Parasite dose, prevalence of infection and local adaptation in a host–parasite system. *Parasitology* **128**, 223–228.
- Overstreet RM and Curran SS** (2004) Defeating diplostomoid dangers in USA catfish aquaculture. *Folia Parasitologica* **51**, 153–165.
- Paperna I** (1991) Diseases caused by parasites in the aquaculture of warm water fish. *Annual Review of Fish Diseases* **1**, 155–194.
- Pérez Ponce de León GPP et al.** (2016) Species delimitation in trematodes using DNA sequences: Middle-American *Clinostomum* as a case study. *Parasitology* **143**, 1773–1789.
- Perpiñán D et al.** (2010) Scoliosis in a tiger salamander (*Ambystoma tigrinum*) associated with encysted digenetic trematodes of the genus *Clinostomum*. *Journal of Wildlife Diseases* **46**, 579–584.
- Pinto HA et al.** (2015) Experimental and molecular study of cercariae of *Clinostomum* sp. (Trematoda: Clinostomidae) from *Biomphalaria* spp. (Mollusca: Planorbidae) in Brazil. *Journal of Parasitology* **101**, 108–113.
- Poulin R and Mouritsen KN** (2003) Large-scale determinants of trematode infections in intertidal gastropods. *Marine Ecology Progress Series* **254**, 187–198.
- R Core Team** (2014) *R: A Language and Environment for Statistical Computing*. Vienna: R Foundation for Statistical Computing. <http://www.R-project.org>.
- Roberts RJ** (2012) *Fish Pathology*. London: John Wiley & Sons.
- Rosser TG et al.** (2016) *Austrodiplostomum* sp., *Bolbophorus* sp. (Digenea: Diplostomidae), and *Clinostomum marginatum* (Digenea: Clinostomidae) metacercariae in inland silverside *Menidia beryllina* from catfish aquaculture ponds, with notes on the infectivity of *Austrodiplostomum* sp. cercariae in channel catfish *Ictalurus punctatus*. *Parasitology Research* **115**, 4365–4378.
- Rosser TG et al.** (2017) *Clinostomum album* n. sp. and *Clinostomum marginatum* (Rudolphi, 1819), parasites of the great egret *Ardea alba* L. from Mississippi, USA. *Systematic Parasitology* **94**, 35–49.
- Rosser TG et al.** (2018) *Clinostomum poteae* n. sp. (Digenea: Clinostomidae), in the trachea of a double-crested cormorant *Phalacrocorax auritus* Lesson, 1831 and molecular data linking the life-cycle stages of *Clinostomum album* Rosser, Alberson, Woodyard, Cunningham, Pote & Griffin, 2017 in Mississippi, USA. *Systematic Parasitology* **95**, 543–566.
- Sandland GJ and Goater CP** (2000) Development and intensity dependence of *Ornithodiplostomum ptychocheilus* metacercariae in fathead minnows (*Pimephales promelas*). *Journal of Parasitology* **86**, 1056–1060.
- Schell SC** (1985) *Handbook of Trematodes of North America North of Mexico*. Moscow, ID: University Press of Idaho.
- Sereno-Uribe AL et al.** (2013) Using mitochondrial and ribosomal DNA sequences to test the taxonomic validity of *Clinostomum complanatum* Rudolphi, 1814 in fish-eating birds and freshwater fishes in Mexico, with the description of a new species. *Parasitology Research* **112**, 2855–2870.
- Sereno-Uribe AL et al.** (2018) Three new species of *Clinostomum* Leidy, 1856 (Trematoda) from Middle American fish-eating birds. *Parasitology Research* **117**, 1–15.
- Sleigh MA** (1991) *Protozoa and Other Protists*. Cambridge: Cambridge University Press.
- Smyth JD and Smyth MM** (1980) *Frogs as Host-Parasite Systems I*. London: Macmillan Press Ltd.
- Szalai AJ and Dick TA** (1988) Helminths of stocked rainbow trout (*Salmo gairdneri*) with special reference to *Clinostomum complanatum*. *Journal of Wildlife Diseases* **24**, 456–460.
- Tkach VV and Pawlowski JW** (1999) A new method of DNA extraction from the ethanol-fixed parasitic worms. *Acta Parasitologica* **44**, 147–148.
- Tkach VV and Snyder SD** (2007) *Aptorchis megacetabulus* n. sp. (Platyhelminthes: Digenea) from the northern long-necked turtle, *Chelodina rugosa* (Pleurodira: Chelidae), in Australia. *Journal of Parasitology* **93**, 404–408.
- Wang ML, Chen HY and Shih HH** (2017) Occurrence and distribution of yellow grub trematode (*Clinostomum complanatum*) infection in Taiwan. *Parasitology Research* **116**, 1761–1771.
- Witenberg G** (1944) What is the cause of the parasitic laryngo-pharyngitis in the Near East ("Halzoun")? *Acta Medica Orientalia* **3**, 191–192.
- Yamaguti S** (1933) Studies on the helminth fauna of Japan. Part I. Trematodes of birds, reptiles and mammals. *Japanese Journal of Zoology* **5**, 1–134.
- Yamashita J** (1938) *Clinostomum complanatum*, a trematode parasite new to man. *Annotationes Zoologicae Japonenses* **17**, 563–566.