



# Meta-analysis on a Decade of Testing Corridor Efficacy: What New Have we Learned?

Julian Resasco<sup>1</sup>

Published online: 21 June 2019

© Springer Nature Switzerland AG 2019

## Abstract

**Purpose of Review** Corridors are widely considered as a strategy to mitigate effects of habitat fragmentation on biodiversity. There are, however, lingering concerns about whether corridors work as intended and whether managing for connectivity in fragmented landscapes is even important for biodiversity conservation. In response, numerous manipulative and natural experiments have been conducted to test the effectiveness of corridors. Gilbert-Norton et al. *Conserv Biol.* 2010;24(3):660–8 (2010) reviewed such studies published between 1985 and 2008 and concluded that corridors are generally effective at increasing inter-patch movement. The authors noted a lack of studies measuring responses at the population and community levels, responses that would better approximate corridor effects on population persistence and aspects of biodiversity. Here I explored what new insights can be gained on corridor effectiveness from studies published in the last decade, particularly with an eye toward insights going beyond effects on inter-patch movement.

**Recent Findings** Following the same selection criteria as Gilbert-Norton et al. *Conserv Biol.* 2010;24(3):660–8 (2010), I reviewed studies published between 2008 and 2018 that tested corridor effectiveness by comparing ecological response variables from patches connected and not connected by corridors. Analysis of effect sizes showed that corridors increase response variables, reinforcing earlier conclusions that corridors function as intended. Whereas the previous review mainly included corridor effects on dispersal, recent research shows support for corridor efficacy at a variety of levels of organization, from individuals to communities.

**Summary** These findings provide further support for the conclusion that efforts spent creating and maintaining corridors are worthwhile for biodiversity conservation.

**Keywords** Corridor experiment · Habitat fragmentation · Conservation · Landscape ecology · Connectivity · Biodiversity

## Introduction

The reduction of connectivity from habitat loss and fragmentation can restrict movement of organisms between sub-populations, which can result in decreased gene flow, local extinctions, and loss of biodiversity

[1, 2]. One method to maintain connectivity is through the conservation or restoration of landscape corridors, strips of habitat that connect otherwise isolated habitat fragments. Despite their intuitive appeal, however, there are long-standing doubts about whether corridors work as intended [3–5]. Assessing the effectiveness of corridors is often challenging due to the difficulty of attaining replication and appropriate controls in landscapes to deal with potentially confounding factors like habitat area and inter-patch distance. Indeed, doubts that took hold about the effectiveness of corridors were in part due to a lack of empirical evidence [4, 6]. In response, a large number of studies have tested corridor function, often controlling for confounding factors.

This article is part of the Topical Collection on *Interface of Landscape Ecology and Conservation Biology*

✉ Julian Resasco  
jresasco@colorado.edu

<sup>1</sup> Department of Ecology & Evolutionary Biology, University of Colorado, UCB 334, Boulder, CO 80309, USA

In a meta-analytic review, Gilbert-Norton et al. [7] synthesized studies (both manipulative and natural experiments) that tested the efficacy of corridors. Reviewing studies from 1985 to 2008, the authors found that: (1) corridors tended to increase movement between habitat patches, (2) the efficacy of corridors varied among taxa, and (3) studies with natural corridors showed greater movement than experimentally manipulated corridors. Given these results, the authors concluded that corridors generally increase movement in fragmented landscapes and therefore corridors are a worthwhile conservation strategy. Gilbert-Norton et al. [7] made significant progress toward addressing the concern about whether corridors function as intended. In addition, concerns of possible negative effects of corridors, such as the spread of disturbance, invasive species, and disease, were addressed in another meta-analysis by Haddad et al. [8]. Yet, Gilbert-Norton et al. [7] also noted that most studies did not test whether inter-patch movement was sufficient to maintain population viability. Moreover, few studies examined population or community level effects. This gap is important because conservation efforts and debates on the effects of landscape structure [9, 10] often focus on the level of populations and communities.

I conducted a meta-analysis on studies testing the effects of corridors to determine whether the past decade of research supports the efficacy of corridors and what new insights can be gained, in particular insights beyond effects on inter-patch movement.

## Methods

I searched for relevant studies on Web of Science (ISI) using the same search terms used by Gilbert-Norton et al. [7]: corridor, movement, effectiveness, connectivity, and habitat connectivity (TS=[corridor AND movement] OR [corridor AND effectiveness] OR [corridor AND connectivity] OR [corridor AND habitat connectivity]). I refined the search results to categories that could include relevant studies: ecology OR environmental sciences OR biodiversity conservation OR zoology OR biology OR behavioral sciences OR soil science OR entomology OR parasitology OR plant sciences OR agronomy. Search results were restricted from 2008 to 2018, allowing me to review the literature since Gilbert-Norton et al. [7]. I also searched the EBSCO Wildlife & Ecology Studies Worldwide database and ProQuest for theses and dissertations.

The Web of Science search resulted in over 1200 results but as in Gilbert-Norton et al. [7], I only included studies if they had replicated corridor and control (unconnected) treatments. Also, as in the previous review, I used the following rules: (1) for studies conducted over multiple time periods, I used the last time period; (2) for studies with multiple types of isolated fragments, I used fragments with a shape that best approximated that of the connected patch; (3) for studies in which data were presented in multiple publications, I used the study that provided the clearest comparison between corridor and control treatments; (4) for studies testing corridors of different widths or lengths, I used the narrowest or longest corridor reasoning that these likely best approximate corridors used in conservation. Similarly, if a study contained treatments with multiple corridors, I used the treatment with the fewest connections. I did not include studies if they were based solely on modeling. I excluded any studies published in 2008 that were already included in Gilbert-Norton et al. [7]. I did not include studies measuring negative interspecific interactions where effects could be ambiguously interpreted as positive for one species and negative for another (e.g., herbivory, [11]). Response variables had to be biotic and included measures like proportion of individuals moved, abundance, and species richness. In cases where multiple closely-related response variables were reported, I selected the variable that I believed to be most meaningful.


For each response variable, I calculated the effect size  $d$ , sometimes referred to as the standardized mean difference or Hedges'  $d$  [12], as in Gilbert-Norton et al. [7]. Values of  $d > 0$  indicate positive effects of corridor treatments compared to unconnected control treatments on ecological response variables. For further interpretation of effect sizes, I use the convention of referring to effect sizes of  $d = 0.2$  as small,  $d = 0.5$  as medium, and  $d = 0.8$  as large [13]. The calculation of  $d$  uses the mean, standard deviation, and sample sizes for corridor-connected and unconnected control groups. In most cases, I was able to extract these values (or data from which these values can be calculated) from the text or figures (using WebPlotDigitizer; <https://automeris.io/WebPlotDigitizer/>) in the paper or by contacting the corresponding author. I was not able to include five studies which did not include the necessary information and for which authors did not respond to requests for data or the values necessary to calculate  $d$ . As in Gilbert-Norton et al. [7], I noted variables of interest for each study. These were: taxon (invertebrates, birds, non-avian vertebrates, and plants); whether studies

controlled for area; whether studies controlled for distance between source and recipient patches; whether studies were manipulative (corridors were created for the experiment) or natural experiments (corridors already present, and if created by humans, not for the purposes of the experiment), and finally whether studies were done at the Savannah River Site Corridor Experiment (South Carolina, USA), a site where many studies meeting the previously stated criteria have been done. I also recorded the level of organization of each response variable as movement, genetic, individual fitness, population, or community.

I analyzed the data using the function “metahdep.HBLM”, hierarchical Bayes linear models allowing for hierarchical dependence, in the R package “metahdep” like in [14], Gilbert-Norton et al. [7]. Gilbert-Norton et al. [7] controlled for sampling dependence which can occur when one control group is compared to multiple experimental groups and hierarchical dependence which can occur when many response variables are measured as a part of a single study. I assumed independence in specifying the model’s dependence structure because all studies analyzed included only a single experimental treatment and most studies had one associated response variable. Differences in effect sizes within studies measuring multiple response variables did not differ from differences among studies measuring single response variables ( $0.86 \pm 0.85$  SD and  $1.03 \pm 0.78$  SD respectively; Welch 2 sample test:  $t = -1.56$ ,  $df = 112.85$ ,  $P = 0.12$ ). I checked that effect sizes conformed to a normal distribution using a Kolmogorov–Smirnov test. I checked that there was not problematic evidence of potential publication bias graphically with a funnel plot and with a normal quantile plot (Wang and Bushman 1998).

## Results

The number of publications from the Web of Science search in the last decade was more than twice those of the previous two decades combined. However, the number of publications that tested the efficacy of corridors in a way that met the criteria for inclusion similar in number to those in Gilbert-Norton et al. [7], 32 here compared to 35 publications for Gilbert-Norton et al. [7]. From these 32 studies, I calculated 56 effect sizes compared to 78 in Gilbert-Norton et al. [7] (Fig. 1a; Appendix Table 1). There were 30 response variables for invertebrates, 13 for non-avian vertebrates (only mammals), 7 for birds, and 6 for plants. There were 11 response variables from 10 papers that examined



**Fig. 1** Effect size ( $d$ ) summaries for studies assessing corridor effectiveness. Values of  $d > 0$  indicate positive effects of corridor treatments on ecological response variables compared to unconnected control treatments. The dashed vertical lines demarcate  $d = 0$ . The top panel (A) is a histogram representing the 56  $d$  values from this meta-analysis. The bottom panel (B) shows estimates of  $d$  from meta-analysis models for different levels of organization as well as for all levels of organization combined (“overall”) in filled gray dots. Error bars represent  $\pm$  SE values from meta-analysis models. Note that the “genetic” category has no error bars because there is a single value of  $d$  associated with it. Colored dots represent corresponding individual  $d$  values with colors representing taxonomic groups. These dots are jittered along the  $y$ -axis to reduce overlap. Vertical gray lines indicate benchmarks of small ( $d = 0.2$ ), medium ( $d = 0.5$ ), and large ( $d = 0.8$ ) effect sizes [13]

movement [16, 20, 23, 27, 28, 32, 37, 39–41], 1 response variable from 1 paper that examined population genetics [45], 4 response variables from 4 papers that examined fitness of individuals [18, 24, 25, 29], 26 response variables from 10 papers that were at the population level [17, 19, 22, 26, 29–31, 42, 43, 46], and 14 responses from 12 papers that were at the community level [15, 21, 26, 31, 33–36, 38, 42, 44, 46] (Appendix Table 1). Manipulated corridors were used to generate 29% of these response variables while natural corridors were used in 71%. Most studies controlled for distance (89%) and area (82%) either experimentally or in site selection. Most responses (88%) did not come from the Savannah River Site Corridor Experiment.

Across all types of response variables, corridors showed an overall positive effect of medium strength ( $d = 0.49$ ,  $P < 0.001$ ; Fig. 1b). For comparison, the mean value of  $d$  for Gilbert-Norton et al. [7] was 0.48. Mean  $d$  values for all levels of organization (movement, genetic, fitness, population, and community) were positive (Fig. 1b). For the movement level,  $d$  was medium-large ( $d = 0.74$ ,  $P < 0.001$ ). For the genetic level,  $d$  was large ( $d = 1.09$ ) but there was only one value of  $d$ . For the fitness level,  $d$  was medium-large ( $d = 0.77$ ,  $P = 0.01$ ). For the population level response variables,  $d$  was medium ( $d = 0.46$ ,  $P < 0.001$ ). For the community level,  $d$  was small ( $d = 0.19$ ,  $P = 0.002$ ).

## Discussion

Results from this meta-analysis reinforce finding that corridors are effective at increasing movement and further extend those conclusions by showing positive effects at a variety of levels of organization, including at the organizational levels of populations and communities. These levels were previously deficiently represented in Gilbert-Norton et al.'s meta-analysis and are more closely aligned with those of conservation goals [47].

While results show that corridors were generally effective, they were not universally effective across studies (e.g., [30, 33, 34] (Fig. 1). Species vary in how they use landscapes and consequently how they respond to fragmentation and corridors [48]. Appropriate tests of corridor effectiveness require appropriate consideration of life history of study organisms and spatial and temporal scale. Most studies reviewed here were surely designed using prior knowledge or intuition about organisms' responses to landscape structure. At the community level, species richness was usually the response variable. Variation in responses to fragmentation among constituent species in species pools may account for the small effect size. That is, species may respond to corridors, but if winners are balanced by losers, then net richness may change little. This is more generally an important consideration in assessments of fragmentation or connectivity effects that consider only species richness [10]. Moreover, timescales could be an important consideration as effects of fragmentation and connectivity can increase with time [2, 49]. Studies reviewed here, as in Gilbert-Norton et al. [7], tended to be short in duration and small in spatial scale compared to typical management activities for biodiversity conservation. Manipulative experiments that are long-term and at the scale of management activities offer unique opportunities to control confounding variables while having high real-world applicability, however, they face strong logistical limitations and so are rare [2, 50, 51].

While manipulative experiments or carefully designed observational studies like those reviewed here offer powerful ways to test the effectiveness of corridors by controlling for confounding factors, other approaches, such as mapping habitat connectivity, simulation studies, animal movement analysis, and behavioral studies can be used to gain system-specific insight as well as contribute to general understanding on effectiveness of corridors. For example, Fletcher et al. [52] broadly reviewed the literature on effects of landscape connectivity and found that, similarly to here, effects of connectivity are frequently observed across levels of organization and are usually positive.

Overall, these findings suggest that to best conserve biodiversity, we need to not only conserve what remains of fragments but conserve and restore their connectivity. This review also points to areas where tests of corridor function were sparse which could guide future studies. As stated previously, studies conducted over large spatial and long temporal scales remain rare but are particularly valuable. While studies at the population and community levels increased in representation, few studies measured responses beyond abundance and species richness to better assess whether corridors reduce extinction risk. There was a surprising lack of studies that tested for effects on gene flow or genetic diversity. There are often mismatches between foci of ecological studies on fragmentation and issues of interest to conservation practitioners, for example, community-based human dimensions, ecosystem services, and interactions of multiple threats to biodiversity [50]. Some of these echo emerging foci for studying fragmentation [2]. Development of research (particularly in co-production with conservation practitioners) in these areas will increase understanding in ecology and in applications for conservation. In closing, current interest in the ecology of corridors is strong; future studies will help reveal mechanisms and contexts for understanding how biodiversity responds to landscape change [9, 10] and allow for increasingly nuanced synthesis.

**Acknowledgments** I thank James Watling and Lenore Fahrig for the invitation to write this paper. I thank Nick Haddad, John Orrock, Doug Levey, Melissa Burt, Lars Brudvig, and Rob Fletcher for valuable feedback. I thank Karen Beard and John Stevens for answering my questions and openly sharing data and code from their meta-analysis. I thank the authors of the reviewed studies who provided additional data. I was supported by a Chancellor's Postdoctoral Fellowship from the University of Colorado while writing this manuscript.

## Compliance with Ethical Standards

**Conflict of Interest** Julian Resasco declares that he has no conflict of interest.

**Human and Animal Rights and Informed Consent** This article does not contain any studies with human or animal subjects performed by any of the authors.

## Appendix

Table 1

| Citation                           | Organisms                                                  | Taxonomic category | Response variable                         | Level of organization | Ne | Nc | Xc         | Se         | Sc       | Sp       | di       |        |
|------------------------------------|------------------------------------------------------------|--------------------|-------------------------------------------|-----------------------|----|----|------------|------------|----------|----------|----------|--------|
| Araujo Calçada et al. (2013) [15]  | Forest herbs                                               | Plant              | Species richness                          | Community             | 8  | 24 | 56.770     | 28.520     | 14.451   | 56.770   | 50.195   | 0.549  |
| Awade and Metzger (2008) [16]      | Variable antshrike ( <i>Thamnophilus caerulexens</i> )     | Bird               | Probability of crossing                   | Movement              | 38 | 38 | 0.813      | 0.500      | 0.403    | 0.509    | 0.459    | 0.674  |
| Awade and Metzger (2008) [16]      | Golden-crowned warbler ( <i>Basileuterus culicivorus</i> ) | Bird               | Probability of crossing                   | Movement              | 38 | 38 | 0.889      | 0.560      | 0.323    | 0.507    | 0.425    | 0.766  |
| Bowler and Benton (2009) [17]      | Mites                                                      | Invertebrate       | Fecundity                                 | Population            | 5  | 5  | 2.720      | 1.650      | 1.160    | 0.570    | 0.914    | 1.057  |
| Crammer et al. (2012) [18]         | Plants                                                     | Plant              | Percent seedset                           | Fitness               | 3  | 4  | 60.587     | 55.488     | 4.376    | 4.317    | 4.341    | 0.989  |
| Evans et al. (2012) [19]           | Grasshoppers                                               | Invertebrate       | Abundance                                 | Population            | 8  | 8  | 43.750     | 40.000     | 6.250    | 6.000    | 6.126    | 0.579  |
| Fletcher et al. (2014) [20]        | Cactus bug ( <i>Chelinidea vittiger</i> )                  | Invertebrate       | Colonization probability                  | Movement              | 10 | 10 | 0.049      | 0.060      | 0.018    | 0.021    | 0.019    | −0.585 |
| Galanes and Thomlinson (2011) [21] | Millipede                                                  | Invertebrate       | Species richness                          | Community             | 6  | 6  | 6.667      | 6.500      | 1.211    | 1.378    | 1.297    | 0.119  |
| Grainger and Gilbert (2017) [22]   | Aphid ( <i>Aphis nerii</i> )                               | Invertebrate       | Max abundance                             | Population            | 7  | 7  | 65.600     | 26.600     | 152.500  | 33.500   | 110.405  | 0.331  |
| Grainger and Gilbert (2017) [22]   | Aphid ( <i>Aphis asclepiadis</i> )                         | Invertebrate       | Max abundance                             | Population            | 7  | 7  | 47.500     | 10.000     | 120.700  | 6.600    | 85.475   | 0.411  |
| Hadley and Betts (2009) [23]       | Green hermit ( <i>Phaethornis guy</i> )                    | Bird               | Homing                                    | Movement              | 10 | 9  | 0.007      | 0.002      | 0.045    | 0.023    | 0.037    | 0.123  |
| Hawn et al. (2018) [24]            | Green Lynx Spiders ( <i>Peucetia viridans</i> )            | Invertebrate       | Lipids                                    | Fitness               | 8  | 8  | 1.380      | 0.930      | 0.460    | 0.160    | 0.344    | 1.235  |
| Herrmann et al. (2017) [25]        | Bumble bee ( <i>Bombus impatiens</i> )                     | Invertebrate       | Abundance (individuals)                   | Fitness               | 8  | 8  | 390.700    | 387.100    | 167.600  | 157.900  | 162.822  | 0.021  |
| Holzschuh et al. (2009) [26]       | Wasps                                                      | Invertebrate       | Species richness                          | Community             | 11 | 9  | 2.636      | 1.667      | 1.502    | 1.500    | 1.501    | 0.619  |
| Holzschuh et al. (2009) [26]       | Wasps                                                      | Invertebrate       | Occupied broodcells                       | Population            | 11 | 9  | 41.273     | 16.000     | 34.523   | 17.937   | 28.375   | 0.853  |
| Ibarra-Macias et al. (2011) [27]   | Birds                                                      | Bird               | Proportion recaptured after translocation | Movement              | 5  | 5  | 0.390      | 0.061      | 0.179    | 0.031    | 0.128    | 2.314  |
| Jesus et al. (2012) [28]           | Plants                                                     | Plant              | Seed rain                                 | Movement              | 3  | 3  | 19,296,000 | 10,164,000 | 1562,000 | 6437,000 | 4683.739 | 1.560  |
| Kormann et al. (2016) [29]         | Heliconia ( <i>Heliconia tortuosa</i> )                    | Plant              | Proportion pollinated flowers             | Fitness               | 13 | 13 | 0.190      | 0.080      | 0.110    | 0.110    | 0.110    | 0.965  |
| Kormann et al. (2016) [29]         | Green hermit ( <i>Phaetornis guy</i> )                     | Bird               | Occupancy probability                     | Population            | 26 | 26 | 0.520      | 0.130      | 0.598    | 1.197    | 0.946    | 0.406  |

Table 1 (continued)

| Citation                          | Organisms                                                               | Taxonomic category | Response variable                                 | Level of organization | Ne | Nc | Xe     | Xc     | Se     | Sc     | Sp     | di            |
|-----------------------------------|-------------------------------------------------------------------------|--------------------|---------------------------------------------------|-----------------------|----|----|--------|--------|--------|--------|--------|---------------|
| Krewenka et al. (2011)            | Solitary bees                                                           | Invertebrate       | Abundance                                         | Population            | 13 | 8  | 1.800  | 1.900  | 0.071  | 0.145  | 0.145  | 0.105 – 0.918 |
| Krewenka et al. (2011) [30]       | Bumble bees                                                             | Invertebrate       | Abundance                                         | Population            | 13 | 8  | 0.880  | 0.750  | 0.031  | 0.045  | 0.037  | 3.365         |
| Kuykendall and Keller (2011) [31] | Rodents                                                                 | Mammal             | Species richness                                  | Community             | 3  | 3  | 7.000  | 8.000  | 1.732  | 0.000  | 1.225  | – 0.653       |
| Kuykendall and Keller (2011) [31] | Western harvest mouse ( <i>Reithrodontomys megalotis</i> )              | Mammal             | Abundance                                         | Population            | 3  | 3  | 75.000 | 42.000 | 46.592 | 38.105 | 42.561 | 0.620         |
| Kuykendall and Keller (2011) [31] | Plains harvest mouse ( <i>Reithrodontomys montanus</i> )                | Mammal             | Abundance                                         | Population            | 3  | 3  | 35.000 | 48.700 | 4.503  | 54.040 | 38.344 | – 0.286       |
| Kuykendall and Keller (2011) [31] | North American deer mouse ( <i>Peromyscus maniculatus</i> )             | Mammal             | Abundance                                         | Population            | 3  | 3  | 25.600 | 17.000 | 20.958 | 12.124 | 17.121 | 0.402         |
| Kuykendall and Keller (2011) [31] | Hispid cotton rat ( <i>Sigmodon hispidus</i> )                          | Mammal             | Abundance                                         | Population            | 3  | 3  | 32.000 | 11.300 | 17.321 | 20.611 | 19.037 | 0.870         |
| Kuykendall and Keller (2011) [31] | Hispid pocket mouse ( <i>Chaetodipus hispidus</i> )                     | Mammal             | Abundance                                         | Population            | 3  | 3  | 13.300 | 14.700 | 11.258 | 12.124 | 11.699 | – 0.096       |
| Kuykendall and Keller (2011) [31] | Northern grasshopper mouse ( <i>Onychomys leucogaster</i> )             | Mammal             | Abundance                                         | Population            | 3  | 3  | 2.700  | 5.700  | 7.967  | 14.030 | 11.409 | – 0.210       |
| Kuykendall and Keller (2011) [31] | Thirteen-lined ground squirrel ( <i>Spermophilus tridecemlineatus</i> ) | Mammal             | Abundance                                         | Population            | 3  | 3  | 1.700  | 5.000  | 2.598  | 10.739 | 7.812  | – 0.338       |
| Kuykendall and Keller (2011) [31] | House mouse ( <i>Mus musculus</i> )                                     | Mammal             | Abundance                                         | Population            | 3  | 3  | 3.000  | 1.300  | 7.448  | 1.039  | 5.317  | 0.256         |
| Kuykendall and Keller (2011) [31] | Ord's kangaroo rat ( <i>Dipodomys ordii</i> )                           | Mammal             | Abundance                                         | Population            | 3  | 3  | 1.000  | 0.700  | 2.944  | 1.905  | 2.480  | 0.097         |
| Leidner and Haddad (2011) [32]    | Crystal skipper ( <i>Atrytonopsis sp1</i> )                             | Invertebrate       | Proportion moved                                  | Movement              | 3  | 6  | 0.320  | 0.098  | 0.185  | 0.120  | 0.142  | 1.388         |
| Orock et al. (2011) [33]          | Arthropods                                                              | Invertebrate       | Species richness                                  | Community             | 7  | 7  | 49.000 | 51.000 | 4.050  | 2.700  | 3.442  | – 0.544       |
| Paolucci et al. (2012) [34]       | Ants                                                                    | Invertebrate       | Species richness                                  | Community             | 5  | 5  | 5.200  | 9.400  | 2.316  | 2.009  | 2.168  | – 1.750       |
| Resasco et al. (2012) [35]        | Red fire ant ( <i>Solenopsis invicta</i> )                              | Invertebrate       | Trophic position                                  | Community             | 10 | 10 | 1.867  | 1.686  | 0.271  | 0.541  | 0.428  | 0.405         |
| Resasco et al. (2014) [36]        | Ants                                                                    | Invertebrate       | Species richness                                  | Community             | 8  | 8  | 10.500 | 13.000 | 4.986  | 4.690  | 4.840  | – 0.488       |
| Riva et al. (2018) [37]           | Arctic fritillary butterfly ( <i>Boloria chariclea</i> )                | Invertebrate       | Probability of persisting in directional movement | Movement              | 6  | 3  | 0.260  | 0.110  | 0.087  | 0.053  | 0.079  | 1.684         |
| Seaman and Schultze (2010) [38]   | Forest-specialist bird                                                  | Bird               | Estimated species richness                        | Community             | 4  | 4  | 13.700 | 9.500  | 4.400  | 9.500  | 7.403  | 0.493         |
| Shimazaki et al. (2016) [39]      | Forest birds                                                            | Bird               | Probability of crossing                           | Movement              | 16 | 12 | 0.798  | 0.472  | 0.228  | 0.243  | 0.235  | 1.349         |
| Van Geert et al. (2010) [40]      | Common primrose ( <i>Primula vulgaris</i> )                             | Plant              | Proportion individuals with pollen dye moved      | Movement              | 13 | 11 | 0.060  | 0.000  | 0.112  | 0.000  | 0.083  | 0.703         |

**Table 1** (continued)

| Citation                          | Organisms                                                | Taxonomic category | Response variable                            | Level of organization | Ne | Nc | Xc         | Se         | Sc       | Sp       | di       |        |
|-----------------------------------|----------------------------------------------------------|--------------------|----------------------------------------------|-----------------------|----|----|------------|------------|----------|----------|----------|--------|
| Van Rossum and Triest (2012) [41] | True oxlip ( <i>Primula elatior</i> )                    | Plant              | Proportion individuals with pollen dye moved | Movement              | 3  | 2  | 0.098      | 0.000      | 0.061    | 0.000    | 0.050    | 1.433  |
| Vergnes et al. (2012) [42]        | Spiders                                                  | Invertebrate       | Species richness                             | Community             | 4  | 4  | 4.940      | 4.090      | 2.260    | 2.280    | 2.270    | 0.326  |
| Vergnes et al. (2012) [42]        | Carabid beetles                                          | Invertebrate       | Species richness                             | Community             | 4  | 4  | 1.580      | 1.160      | 2.440    | 2.480    | 2.460    | 0.148  |
| Vergnes et al. (2012) [42]        | Staphylinid beetles                                      | Invertebrate       | Species richness                             | Community             | 4  | 4  | 3.320      | 1.290      | 2.320    | 2.420    | 2.371    | 0.745  |
| Vergnes et al. (2012) [42]        | Spiders                                                  | Invertebrate       | Abundance                                    | Population            | 4  | 4  | 10.960     | 6.500      | 2.300    | 2.320    | 2.310    | 1.679  |
| Vergnes et al. (2012) [42]        | Carabid beetles                                          | Invertebrate       | Abundance                                    | Population            | 4  | 4  | 2.190      | 1.840      | 2.840    | 2.860    | 2.850    | 0.107  |
| Vergnes et al. (2012) [42]        | Staphylinid beetles                                      | Invertebrate       | Abundance                                    | Population            | 4  | 4  | 11.160     | 3.870      | 2.480    | 2.520    | 2.500    | 2.536  |
| Vergnes et al. (2013) [43]        | Greater white-toothed shrew ( <i>Crocidura russula</i> ) | Mammal             | Occupancy                                    | Population            | 4  | 4  | 1.750      | 1.670      | 0.850    | 0.630    | 0.748    | 0.093  |
| Vergnes et al. (2013) [43]        | Crowned shrew ( <i>Sorex coronatus</i> )                 | Mammal             | Occupancy                                    | Population            | 4  | 4  | 1.500      | 0.500      | 1.500    | 0.500    | 1.118    | 0.778  |
| Vergnes et al. (2013) [43]        | Eurasian pygmy shrew ( <i>Sorex minutus</i> )            | Mammal             | Occupancy                                    | Population            | 4  | 4  | 0.750      | 2.500      | 0.750    | 2.170    | 1.623    | −0.937 |
| Viljur et al. (2018) [44]         | Open habitat Butterflies                                 | Invertebrate       | Species richness                             | Community             | 18 | 18 | 13.100     | 11.800     | 3.100    | 2.700    | 2.907    | 0.437  |
| Wells et al. (2009) [45]          | Buckeye butterfly ( <i>Junonia coenia</i> )              | Invertebrate       | Heterozygosity (observed)                    | Genetic               | 6  | 6  | 0.193      | 0.155      | 0.043    | 0.015    | 0.032    | 1.086  |
| Zaitsev et al. (2014) [46]        | Soil Macrofauna                                          | Invertebrate       | Taxon richness                               | Community             | 3  | 3  | 18.000     | 13.000     | 1.039    | 1.559    | 1.325    | 3.019  |
| Zaitsev et al. (2014) [46]        | Soil Macrofauna                                          | Invertebrate       | Abundance                                    | Population            | 3  | 3  | 2573.000   | 2040.000   | 478.046  | 1290.378 | 973.037  | 0.438  |
| Zaitsev et al. (2014) [46]        | Oribatid mites                                           | Invertebrate       | Abundance                                    | Population            | 3  | 3  | 30,250.000 | 19,533.000 | 7996.879 | 6283.880 | 7191.565 | 1.192  |
| Zaitsev et al. (2014) [46]        | Mesostigmatid mites                                      | Invertebrate       | Abundance                                    | Population            | 3  | 3  | 2633.000   | 2067.000   | 1357.928 | 923.183  | 1161.085 | 0.390  |

## References

- Hilty JA, Lidicker WZ, Merenlender AM. Corridor ecology : the science and practice of linking landscapes for biodiversity conservation. Washington, DC: Island Press; 2006.
- Haddad NM, Brudvig LA, Clobert J, Davies KF, Gonzalez A, Holt RD, et al. Habitat fragmentation and its lasting impact on Earth's ecosystems. *Sci Adv*. 2015;1(2):E1500052.
- Simberloff D, Cox J. Consequences and costs of conservation corridors. *Conserv Biol*. 1987;1:63–71.
- Simberloff D, Farr JA, Cox J, Mehlman DW. Movement corridors: conservation bargains or poor investments? *Conserv Biol*. 1992;6(4):493–504.
- Jain A, Chong KY, Chua MAH, Clements GR. Moving away from paper corridors in Southeast Asia. *Conserv Biol*. 2014;28(4):889–91.
- Beier P, Noss RF. Do habitat corridors provide connectivity? *Conserv Biol*. 1998;12(6):1241–52.
- Gilbert-Norton L, Wilson R, Stevens JR, Beard KH. A meta-analytic review of corridor effectiveness. *Conserv Biol*. 2010;24(3):660–8.
- Haddad NM, Brudvig LA, Damschen EI, Evans DM, Johnson BL, Levey DJ et al. A review of potential negative ecological effects of corridors. *Conserv Biol*. 2014;28(5):1178–87.
- Fahrig L, Arroyo-Rodríguez V, Bennett JR, Boucher-Lalonde V, Cazetta E, Currie DJ, et al. Is habitat fragmentation bad for biodiversity? *Biol Conserv*. 2019;230:179–86.
- Fletcher RJ, Didham RK, Banks-Leite C, Barlow J, Ewers RM, Rosindell J, et al. Is habitat fragmentation good for biodiversity? *Biol Conserv*. 2018;226:9–15.
- Levey DJ, Caughlin TT, Brudvig LA, Haddad NM, Damschen EI, Tewksbury JJ, et al. Disentangling fragmentation effects on herbivory in understory plants of longleaf pine savanna. *Ecology*. 2016;97(9):2248–58.
- Gurevitch J, Curtis PS, Jones MH. Meta-analysis in ecology. *Adv Ecol Res*. 2001;32:199–247.
- Cohen J. Statistical power analysis for the behavioral sciences. 2nd ed. Hillsdale, N.J.: L. Erlbaum Associates; 1988.
- Stevens JR, Metahdep NG. Hierarchical dependence in meta-analysis. In: R package version 1.34.0. ed; 2011.
- Araujo Calçada E, Closset-Kopp D, Gallet-Moron E, Lenoir J, Reve M, Henny M, et al. Streams are efficient corridors for plant species in forest metacommunities. *J Appl Ecol*. 2013;50(5):1152–60.
- Awade M, Metzger JP. Using gap-crossing capacity to evaluate functional connectivity of two Atlantic rainforest birds and their response to fragmentation. *Austral Ecol*. 2008;33(7):863–71.
- Bowler DE, Benton TG. Impact of dispersal on population growth: the role of inter-patch distance. *Oikos*. 2009;118(3):403–12.
- Cranmer L, McCollin D, Ollerton J. Landscape structure influences pollinator movements and directly affects plant reproductive success. *Oikos*. 2012;121(4):562–8.
- Evans DM, Turley NE, Levey DJ, Tewksbury JJ. Habitat patch shape, not corridors, determines herbivory and fruit production of an annual plant. *Ecology*. 2012;93(5):1016–25.
- Fletcher RJ, Acevedo MA, Robertson EP. The matrix alters the role of path redundancy on patch colonization rates. *Ecology*. 2014;95(6):1444–50.
- Galanes IT, Thomlinson JR. Soil millipede diversity in tropical forest patches and its relation to landscape structure in northeastern Puerto Rico. *Biodivers Conserv*. 2011;20(13):2967–80.
- Grainger TN, Gilbert B. Multi-scale responses to warming in an experimental insect metacommunity. *Glob Chang Biol*. 2017;23(12):5151–63.
- Hadley AS, Betts MG. Tropical deforestation alters hummingbird movement patterns. *Biol Lett*. 2009;5(2):207–10.
- Hawn CL, Herrmann JD, Griffin SR, Haddad NM. Connectivity increases trophic subsidies in fragmented landscapes. *Ecol Lett*. 2018;21(11):1620–8.
- Herrmann JD, Haddad NM, Levey DJ. Testing the relative importance of local resources and landscape connectivity on *Bombus impatiens* (hymenoptera, Apidae) colonies. *Apidologie*. 2017;48(4):545–55.
- Holzschuh A, Steffan-Dewenter I, Tschamtk T. Grass strip corridors in agricultural landscapes enhance nest-site colonization by solitary wasps. *Ecol Appl*. 2009;19(1):123–32.
- Ibarra-Macias A, Robinson WD, Gaines MS. Forest corridors facilitate movement of tropical forest birds after experimental translocations in a fragmented Neotropical landscape in Mexico. *J Trop Ecol*. 2011;27:547–56.
- Jesus FM, Pivello VR, Meirelles ST, Franco GADC, Metzger JP. The importance of landscape structure for seed dispersal in rain forest fragments. *J Veg Sci*. 2012;23(6):1126–36.
- Kormann U, Scherber C, Tschamtk T, Klein N, Larbig M, Valente JJ, et al. Corridors restore animal-mediated pollination in fragmented tropical forest landscapes. *P Roy Soc B-Biol Sci*. 2016;283(1823).
- Krewenka KM, Holzschuh A, Tschamtk T, Dormann CF. Landscape elements as potential barriers and corridors for bees, wasps and parasitoids. *Biol Conserv*. 2011;144(6):1816–25.
- Kuykendall MT, Keller GS. Impacts of roads and corridors on abundance and movement of small mammals on the Llano Estacado of Texas. *Southwest Nat*. 2011;56(1):9–16.
- Leidner AK, Haddad NM. Combining measures of dispersal to identify conservation strategies in fragmented landscapes. *Conserv Biol*. 2011;25(5):1022–31.
- Orrock JL, Curler GR, Danielson BJ, Coyle DR. Large-scale experimental landscapes reveal distinctive effects of patch shape and connectivity on arthropod communities. *Landsc Ecol*. 2011;26(10):1361–72.
- Paolucci LN, Solar RRC, Sobrinho TG, Sperber CF, Schoederer JH. How does small-scale fragmentation affect litter-dwelling ants? The role of isolation. *Biodivers Conserv*. 2012;21(12):3095–105.
- Resasco J, Levey DJ, Damschen EI. Habitat corridors alter relative trophic position of fire ants. *Ecosphere*. 2012;3(11).
- Resasco J, Haddad NM, Orrock JL, Shoemaker D, Brudvig TA, Damschen EI, et al. Landscape corridors can increase invasion by an exotic species and reduce diversity of native species. *Ecology*. 2014;95(8):2033–9.
- Riva F, Acorn JH, Nielsen SE. Narrow anthropogenic corridors direct the movement of a generalist boreal butterfly. *Biol Lett*. 2018;14(2).
- Seaman BS, Schulze CH. The importance of gallery forests in the tropical lowlands of Costa Rica for understory forest birds. *Biol Conserv*. 2010;143(2):391–8.
- Shimazaki A, Yamaura Y, Senzaki M, Yabuhara Y, Akasaka T, Nakamura F. Urban permeability for birds: an approach combining mobbing-call experiments and circuit theory. *Urban Urban Gree*. 2016;19:167–75.
- Van Geert A, Van Rossum F, Triest L. Do linear landscape elements in farmland act as biological corridors for pollen dispersal? *J Ecol*. 2010;98(1):178–87.
- Van Rossum F, Triest L. Stepping-stone populations in linear landscape elements increase pollen dispersal between urban forest fragments. *Plant Ecol Evol*. 2012;145(3):332–40.
- Vergnes A, Le Viol I, Clergeau P. Green corridors in urban landscapes affect the arthropod communities of domestic gardens. *Biol Conserv*. 2012;145(1):171–8.

43. Vergnes A, Kerbiriou C, Clergeau P. Ecological corridors also operate in an urban matrix: a test case with garden shrews. *Urban Ecosyst*. 2013;16(3):511–25.
44. Viljur ML, Teder T. Disperse or die: colonisation of transient open habitats in production forests is only weakly dispersal-limited in butterflies. *Biol Conserv*. 2018;218:32–40.
45. Wells CN, Williams RS, Walker GL, Haddad NM. Effects of corridors on genetics of a butterfly in a landscape experiment. *Southeast Nat*. 2009;8(4):709–22.
46. Zaitsev AS, Gongalsky KB, Persson T, Bengtsson J. Connectivity of litter islands remaining after a fire and unburnt forest determines the recovery of soil fauna. *Appl Soil Ecol*. 2014;83:101–8.
47. Gregory AJ, Beier P. Response variables for evaluation of the effectiveness of conservation corridors. *Conserv Biol*. 2014;28(3):689–95.
48. Brudvig LA, Leroux SJ, Albert CH, Bruna EM, Davies KF, Ewers RM, et al. Evaluating conceptual models of landscape change. *Ecography*. 2017;40(1):74–84.
49. Damschen EI, Brudvig LA, Burt MA, Fletcher RJ, Haddad NM, Levey DJ et al. Landscape connectivity promotes colonization credits and reduces extinction debt over decades. in prep.
50. Resasco J, Bruna EM, Haddad NM, Banks-Leite C, Margules CR. The contribution of theory and experiments to conservation in fragmented landscapes. *Ecography*. 2017;40(1):109–18.
51. Debinski DM, Holt RD. A survey and overview of habitat fragmentation experiments. *Conserv Biol*. 2000;14(2):342–55.
52. Fletcher RJ, Burrell NS, Reichert BE, Vasudev D, Austin JD. Divergent perspectives on landscape connectivity reveal consistent effects from genes to communities. *Curr Landscape Ecol Rep*. 2016;1(2):67–79.

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.