

Youth Screen Media Habits and Sleep

Sleep-Friendly Screen Behavior Recommendations for Clinicians, Educators, and Parents



Lauren Hale, PhD^{a,*}, Gregory W. Kirschen, PhD^b,
Monique K. LeBourgeois, PhD^c, Michael Gradisar, PhD^d,
Michelle M. Garrison, PhD^{e,f}, Hawley Montgomery-Downs, PhD^g,
Howard Kirschen, MD^h, Susan M. McHale, PhDⁱ,
Anne-Marie Chang, PhD^{j,k}, Orfeu M. Buxton, PhD^{l,m,n}

KEYWORDS

• Youth • Screen media habits • Sleep • Screen behavior recommendations

Disclosures: Authors on this paper were supported in part by the Eunice Kennedy Shriver National Institute of Child Health and Human Development (NICHD) of the National Institutes of Health (NIH) under award numbers R01HD073352 (supporting Dr L. Hale, Dr A.M. Chang, G.W. Kirschen, and Dr O.M. Buxton), R01HD087707 (supporting Dr M.K. LeBourgeois), and R01HD071937 (supporting Dr M.M. Garrison). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. Outside of the current work, Dr O.M. Buxton received subcontracts from Mobile Sleep Technologies for National Science Foundation award 1622766 and NIH/National Institute on Aging (NIA) R43AG056250. Dr M. Gradisar has received consultancies from Johnson & Johnson, the Australian Psychological Society, the National Health & Medical Research Council, Access Macquarie, and Little Brown Book Company.

^a Program in Public Health, Department of Family, Population, and Preventive Medicine, Stony Brook Medicine, HSC Level 3, Room 071, Stony Brook, NY 11794-8338, USA; ^b Medical Scientist Training Program, Stony Brook Medicine, HSC Level 3, Room 071, Stony Brook, NY 11794-8338, USA; ^c Department of Integrative Physiology, University of Colorado Boulder, Boulder, CO 80309-0354, USA; ^d Department of Psychology, Flinders University, GPO Box 2100, Adelaide 5001, South Australia; ^e Division of Child and Adolescent Psychiatry, University of Washington School of Medicine, 4333 Brooklyn Avenue NE, Seattle, WA 98195-9455, USA; ^f Department of Health Services, University of Washington School of Public Health, Seattle, WA, USA; ^g Department of Psychology, West Virginia University, PO Box 6040, 53 Campus Drive, 1124 LSB, Morgantown, WV 26506-6040, USA; ^h Child, Adolescent, Adult Psychiatry and Psychotherapy Private Practice, 366 N Broadway Street 210, Jericho, NY 11753, USA; ⁱ Department of Human Development and Family Studies, The Pennsylvania State University, 114 Henderson, University Park, PA 16802, USA; ^j Department of Biobehavioral Health, The Pennsylvania State University, Biobehavioral Health Building, University Park, PA 16802, USA; ^k College of Nursing, The Pennsylvania State University, Nursing Sciences Building, University Park, PA 16802, USA; ^l Division of Sleep Medicine, Harvard Medical School, 221 Longwood Avenue, Boston, MA 02115, USA; ^m Division of Sleep and Circadian Disorders, Departments of Medicine and Neurology, Sleep Health Institute, Brigham and Women's Hospital, 221 Longwood Avenue, Boston, MA 02115, USA; ⁿ Department of Social and Behavioral Sciences, Harvard Chan School of Public Health, 677 Huntington Avenue, Boston, MA 02115, USA

* Corresponding author.

E-mail addresses: Lauren.Hale@stonybrook.edu; Lauren.Hale@stonybrookmedicine.edu

Child Adolesc Psychiatric Clin N Am 27 (2018) 229–245

<https://doi.org/10.1016/j.chc.2017.11.014>

childpsych.theclinics.com

1056-4993/18/© 2017 Elsevier Inc. All rights reserved.

KEY POINTS

- Use of screen media by youth is associated with shorter total sleep time, delayed sleep onset, shorter sleep duration, later bedtime, and poorer sleep quality.
- Mechanisms underlying the relationship between screen media habits and sleep outcomes include displacement of sleep time spent, psychological stimulation from content, and (3) alerting and circadian effects of exposure to light from screens.
- Clinicians, educators, and parents should prioritize the need of youth to get sufficient sleep by maintaining regular and consistent bedtime routines.
- There is a need for more basic, translational, and clinical research examining the effects of screen media on sleep and related health and health behavior consequences.

Abbreviations

FERRET	Food, Emotions, Routine, Restrict, Environment and Timing
IVGA	Internet and video game addiction
NSF	National Sleep Foundation

INTRODUCTION

The widespread use of portable electronic devices and the normalization of screen media devices in the bedroom is accompanied by a high prevalence of insufficient sleep, affecting a majority of adolescents, and 30% of toddlers, preschoolers, and school-age children.^{1–6} Three-fourths of American children and adolescents report the presence of at least 1 screen media device in their bedroom, with roughly 60% reporting regular use of these devices during the hour before bedtime.^{7,8}

Parents, educators, and clinicians express concern about whether excessive use of screen media among young people affects sleep and well-being. In this article, we provide an overview of the current science on screens and sleep, with a focus on recommendations to reduce the potentially problematic influence of screen time on pediatric sleep. We then review how impaired sleep in pediatric populations may lead to a range of adverse behaviors, physical health problems, and well-being outcomes. We begin with a summary of the 2 consensus statements on child and adolescent sleep needs. Then, we summarize the range of screen habits among youth, focusing on screen habits at bedtime. Next, we review current literature on evidence of the effects of youth screen habits on sleep, and the mechanisms by which screen habits may impact sleep. We conclude with evidence-based strategies to improve sleep through sleep-friendly screen behavior recommendations and other take-home messages for families and practitioners.

SLEEP REQUIREMENTS FOR CHILDREN AND ADOLESCENTS

Two independent sleep associations—the National Sleep Foundation (NSF) and American Academy of Sleep Medicine—each convened teams of sleep researchers and other experts to establish consensus to guide health care providers and the public about sleep duration requirements across the lifespan, based on the best available evidence. Both groups used a modified RAND/UCLA Appropriateness Method⁹ to arrive at their recommendations. For the pediatric population, the NSF panel

recommended that newborns (0–3 months) obtain 14 to 17 hours of sleep daily, infants (4–11 months) obtain 12 to 15 hours, toddlers (1–2 years) obtain 11 to 14 hours, preschoolers (3–5 years) obtain 10 to 13 hours, school-aged children (6–13 years) obtain 9 to 11 hours, and teenagers (14–17 years) obtain 8 to 10 hours.¹⁰ The American Academy of Sleep Medicine recommendations were identical to those of the NSF, except the former suggested that infants aged 4 to 12 months obtain 12 to 16 hours of sleep per day (including naps) and children aged 6 to 12 obtain 9 to 12 hours.¹¹

Consequences of Insufficient Sleep for Cognitive, Psychological, and Physical Well-Being

Sleep problems in early life predict a greater likelihood of later development of psychopathology in childhood and adolescence.¹² In a large study (N = 32,662), short sleep duration (≤ 10 hours per night by maternal report) and nocturnal awakenings (≥ 3 per night) in toddlers were associated with the development of behavioral and emotional problems at age 5.¹³ Sleep problems at age 4 have been found to predict a greater incidence of behavioral and emotional problems emerging by mid-adolescence.¹⁴ In experimental studies, toddler napping seems to be important for self-regulation. A challenging puzzle task after 5 days of regular napping (compared with not napping) elicited fewer perseverative behaviors, and less negative self-appraisals.¹⁵ Imposing sleep restriction on healthy teens rapidly degrades mood and emotion regulation,¹⁶ as with adults. A large cross-sectional study of adolescents identified associations between short sleep duration and emotional problems, peer conflict, and suicidal ideation.¹⁷ In a metaanalysis of longitudinal and intervention studies, adolescent sleep problems seem to precede the emergence of depression.¹⁸ Although more work on the modifiable aspects of sleep duration, sleep quality, and regularity of sleep timing are needed, the current literature suggests that each of these factors is important for psychological health. These relationships may be bidirectional, because sleep and psychological problems influence one another throughout development,¹⁹ suggesting a resonance phenomenon or a vicious cycle that may be exacerbated by excessive screen media habits. Williams and colleagues²⁰ have proposed a developmental cascade model, supported by longitudinal data on children, to explain the reciprocal interactions of sleep and emotion and attention self-regulation.

Beyond emotional and behavioral problems, a metaanalysis of cross-sectional studies revealed that short sleep duration was reliably associated with weight gain, adiposity, and obesity risk in both children and adults.⁴ Infants obtaining less than 12 hours of sleep per day, measured by maternal report, were more likely to be overweight at age 3 and have higher levels of adiposity.²¹ In a longitudinal study of children age 1 to 7 years, sleep duration was associated with greater increases in body mass index, fat mass, and waist-to-hip ratio.²² In a controlled, laboratory-based experimental study, 3-year-old children exposed to acute sleep restriction (skipping a nap and bedtime delayed by approximately 2.3 hours) consumed greater amounts of carbohydrate and fat and more total calories,²³ which may explain the increased risk of obesity caused by insufficient sleep. A recent review identified sleep as among the socioeconomic, family, environmental, and behavioral factors contributing to childhood obesity.²⁴

In a longitudinal study of children in the third grade, bedtimes after 8 PM were associated with a greater odds of increased adiposity measured in the sixth grade. As with children, insufficient sleep in adolescence has been shown to promote dietary behaviors that lead to obesity,²⁵ with additional effects mediated via decreased physical activity and neuroendocrine changes that bolster fat storage. More evidence is needed, particularly in interventions that counteract the effects of screen media consumption leading to insufficient quantity or inadequate quality sleep, that in turn influence weight

gain in childhood. However, the “weight” of available evidence suggests that obesity may be a sleep loss-related outcome of excessive screen media consumption.

WHAT ARE THE BEDTIME SCREEN HABITS OF INFANTS, CHILDREN, AND ADOLESCENTS?

Although the scientific and clinical communities continue to express concern regarding the negative impacts of screen media on sleep, electronics in the bedroom and screen time use around bedtime remain common among youth.^{26–28} A large-scale, nationally representative sample of American parents in 2013²⁶ revealed that about one-third of young children (36%) had televisions in their bedrooms, including 16% of children under 2 years of age, 37% of 2- to 4-year-olds, and 45% of 5- to 8-year-olds. Additionally, among parents of infants and young children who allowed a bedroom television, 22% did so to help their child fall asleep, 14% did so to get their child to fall asleep in his or her own room, and 4% did so because their child slept in a family room containing a television.²⁶ The presence of a television in the bedroom varies by cultural, socioeconomic, and structural factors. Data from a 2016 nationally representative study of more than 2600 US youth indicated that 47% of tweens and 57% of teens have televisions in their bedrooms.²⁷ Media devices in the bedroom were more common in lower income youth (68% vs 37%). The authors speculated that the higher rate of media in the bedrooms of lower income tweens and teens may be the result of more frequent room sharing, sleeping in a multipurposed room, or differing family preferences. Parent-reported data from the 2014 *Sleep in America* poll by the NSF revealed that 75% of youth keep at least 1 type of electronic device in their bedroom. The poll also found that the 28% of school-aged children and 57% of teenagers who leave an electronic device on in their bedroom after bedtime obtained less total sleep and had lower sleep quality.⁷

About 50% of parents endorse the belief that watching television helps their infant, toddler, or preschooler “wind down” in the evening.²⁸ Likewise, multiple studies documented that adolescents report using media at bedtime to “help” them fall asleep.²⁹ There is a current programming trend toward developing calming shows and apps aimed at helping children relax before bedtime and transition to sleep. To our knowledge, no published research exists on the effectiveness of such content. Such evidence should be a minimum requirement for these approaches to be recommended.

Importantly, over the past decade, the landscape of screen media devices has changed markedly, dramatically altering children and adolescents engage with their environment. Data from large national US samples in 2011 and 2013 show an increase in ownership of tablet devices in young children (age 0–8) from 8% to 40% and an increase in access to mobile “smart” devices at home from 52% to 75%.²⁶ Another report indicated that, from 2011 to 2013, the percentage of children under 2 years of age who had used a mobile device increased from 10% to 38%.²⁶

SCREEN MEDIA, ESPECIALLY NEAR BEDTIME, IS ADVERSELY ASSOCIATED WITH SLEEP TIME AND QUANTITY

In relation to a growing interest in the association between screen time and sleep patterns, data from more than 60 associated studies have been examined in 2 systematic literature reviews^{3,30} and 2 related metaanalyses.^{31,32} In the time since those articles were published, at least a dozen more studies have surfaced (eg, see^{6,8,33–42}), from a wide range of cultural contexts including Thailand,³⁷ Saudi Arabia,⁴⁰ and Norway.⁸

The vast majority of these studies indicate that the extent of screen time among children and adolescents is associated with delayed bedtime and shorter total sleep time.^{43–45} Several studies also found associations between screen time and reduced

sleep quality,^{46,47} longer sleep onset latency,⁴⁸ and increased daytime tiredness.^{49,50} One recent study found that electronic media use accounted for 30% of all variance in adolescent sleep efficiency, as measured by actigraphy.³⁴ Most studies examined total daily screen time as a predictor, but even greater effects on sleep have been documented in evening media use in the bedroom (ie, in the 1–2 hours before bedtime^{51,52}) and in use of violent media at any time.^{53,54}

One metaanalysis investigated the association between portable screen-based media devices and sleep outcomes.³² Merging results from 20 studies and more than 125,000 youth, the authors consistently found that bedtime media usage is associated with insufficient sleep duration (odds ratio, 2.17; $P < .001$), poor sleep quality (odds ratio, 1.46; $P < .01$), and excessive daytime sleepiness (odds ratio, 2.72; $P < .01$). This metaanalysis found that the mere presence of a portable screen-based media device in the bedroom has adverse associations with sleep outcomes. In 1 study of 600 preschoolers, those with a television in their bedroom watched twice as much evening television (about 28 minutes vs 13 minutes, respectively) and watched more shows with violent, scary, or “mature” content (29% vs 13%).⁵³ Most related studies focused on typically developing children, but results of studies on children with attention deficit hyperactivity disorder and autism spectrum disorders show similar patterns. One such study found that exposure to violent programming within 30 minutes before bedtime was associated with longer sleep latency and shorter sleep duration.⁵⁴

Negative associations with sleep have been found with use of a range of screen media devices, including televisions, computers, video games, and mobile devices such as smartphones and tablets.^{29,48,51,52,55–64} However, there are mixed results regarding whether the type, size, or interactivity level of the screen affects sleep outcomes. In a study of 2048 children in grades 4 and 7, having either a television or small screen device near where they sleep—as well as more daily screen time—was associated with obtaining less sleep.⁶ The increase in interactive media options may increase the impact of media use on sleep; some studies have found that interactive screen media use (eg, video games and mobile devices) may have a greater impact on sleep than passive use, such as watching television.^{6,45,65–70} One recent metaanalysis found no association between television watching and sleep duration,³¹ but did find that computer use is associated with a shorter total sleep duration. Another study shows that use of interactive screen media increases the odds of nighttime awakenings and daytime tiredness,⁴¹ but other studies examining the effects of video game use on sleep have shown more modest effects.^{67,71} One study compared 1 hour of bedtime tablet use with 3 different lighting profiles and found minimal differences in the impact on sleep and next-day functioning.⁷² However, longer durations of bright screen use (between 1.5 and 5.0 hours) have been shown to increase alertness before sleep.^{73–76}

MECHANISMS THROUGH WHICH SCREENS AFFECT SLEEP

Because many of the existing studies are observational and cross-sectional, causality is difficult to discern. Several potential mechanisms, along with supporting evidence, are briefly discussed, including time displacement, psychological stimulation from content, and the alerting and circadian effects of light.

Time Displacement of Sleep

Screen media use can lead to behavioral bedtime delay, as children or adolescents postpone bedtime to prolong screen entertainment.^{65,77,78} Using a screen-based device displaces time that would otherwise have been spent sleeping. This time

displacement mechanism is particularly powerful when screens are used at night, when sleep is most likely the activity being directly offset.

Psychological Stimulation from Media Content Disrupting Sleep

Research has found mixed results regarding the effect of screen media use on psychological, emotional, and/or physiologic arousal, but this relationship likely mediates some effect of screen media use on sleep.^{79–82} Video games, particularly violent games, are often thrilling for enthusiasts, typically simulating life-or-death experiences requiring players' full attention to succeed. Violent video game play before bed increase arousal compared with nonviolent gameplay. However, arousal is a likely mediator of sleep problems even in nonviolent media use.⁴⁶ In another controlled experiment, playing an "exciting" video game was associated with increased heart rate, slightly delayed sleep-onset, and decreased REM sleep, further suggesting that the effect of screen use on sleep is mediated via arousal.⁶⁰

Effects of Light-Emitting Screens on Child and Adolescent Sleep

Exposure to the light emitted by screens in the evening hours before and/or during bedtime is another likely mechanism by which use of screen media negatively impacts sleep. Screen-based light may affect sleep via several pathways:

1. Increasing arousal and reducing sleepiness at bedtime,
2. Disrupting sleep architecture as assessed by polysomnographic recording, and
3. Delaying the circadian rhythm and subsequently postponing sleep-onset, which results in shortened sleep duration unless wake time is also delayed.

Studies of young adults demonstrated that evening use of light-emitting devices increases alertness and decreases sleepiness before bedtime, as determined by cognitive performance, self-reported scales, and waking electroencephalographic measures.^{73,75,83} These studies also showed that this light suppresses blood levels of the sleep-promoting hormone melatonin, which normally increases in the hours before bedtime.^{73,75,76,83} Similar results were found in an experimental study of adolescents, in which exposure to LED screens before bedtime decreased self-reported sleepiness and melatonin levels.⁷⁴ Exposure to nonscreen artificial light in the evening also increases alertness (quelling underlying sleep drive) and suppresses melatonin levels, delaying sleep onset. The light emitted from screen media devices includes greater short-wavelength light in the blue light range, which has been shown to induce stronger melatonin-suppressing responses.⁸⁴ In fact, in the aforementioned study of adolescents,⁷⁴ both subjective alertness and melatonin suppression were significantly attenuated when study participants wore glasses that filtered out short-wavelength light.

Few studies have directly measured the effects of light from screens on polysomnographic measures of sleep.^{72,74,75} One such study in young adults found that exposure to light-emitting devices before bedtime caused a phase delay of melatonin release and modestly increased the time to fall asleep and reduced the duration of REM sleep.⁷⁵ It is impossible to determine whether longer sleep latency or reduced REM sleep duration was due to decreased sleepiness before bedtime, suppression of melatonin, a phase delay of the circadian clock, or a combination of these factors. Other published reports, including a study of adolescents,⁷² found no significant changes on subsequent polysomnography with evening exposure to light-emitting devices.^{72,74}

Although limited research documents the effects of light from digital media on sleep, the results are particularly applicable to youth, who may be more sensitive to light than adults.⁸⁵ One study⁸⁶ found that the magnitude of melatonin suppression induced by moderately bright indoor light levels in the evening was twice as much in primary

school children as in adults. This difference could be related to pupil diameter, which is significantly larger in children than adults. Another study⁸⁷ found that children showed significantly greater melatonin suppression compared with adolescents in response to varying degrees of evening light exposure. Taken together, these results suggest that the light from screen media use around bedtime adversely affects sleep, particularly in younger children. Furthermore, these findings indicate the need for further research on the effect of screen media devices on sleep, especially in youth.

INTERNET AND VIDEO GAME ADDICTION AND SLEEP

There is growing concern that technology habits can become uncontrollable and excessive to the point of interference with normal daily functioning. For example, the *Diagnostic and Statistical Manual of Mental Disorders*, 5th edition, recognizes 1 type of technology addiction, Internet gaming disorder, as a “condition for further study.”⁸⁸ More broadly, the concept of Internet and video game addiction (IVGA), has gained traction in the psychological and psychiatric communities, due in part to the disorder’s deleterious effects on sleep. IVGA is classified as an inability of Internet users to limit excessive Internet use, with ensuing psychosocial dysfunction.⁸⁹ A recent systematic literature review and other studies show a particularly strong association between use of massively multiplayer online role-playing games and poor sleep quality.^{90–92} IVGA has been associated with subjective insomnia and insufficient sleep among afflicted adolescents.⁹⁰ Existing literature on this association remains limited and causality is unproven, but there seems to be a strong association between IVGA and sleep problems.

MODERATING FACTORS

New evidence suggests some young people are particularly susceptible to the ill effects of screen media on sleep quality. As early as 2009, researchers documented individual differences in teenagers’ heart rate variability when playing violent video games.⁸² A subsequent study⁴⁶ demonstrated that adolescents’ level of gaming experience moderated the effect between their video game use, heart rate variability, and sleep. Inexperienced gamers reported poorer sleep after playing a violent video game, in contrast with experienced gamers, who reported poorer sleep after a nonviolent video game. Physiologic trait differences can also amplify technology exposure effects by directly delaying bedtimes.⁵² An experimental laboratory study by Reynolds and colleagues⁷⁸ revealed that adolescents who perceived fewer consequences from risk taking were more likely to end daily video gaming sessions at a later time, thus delaying sleep onset, compared with their peers. Smith and colleagues⁹³ found that adolescents who self-reported higher trait flow (the ability to easily immerse oneself into an activity) were also more likely to delay bedtime via extended periods of evening gaming. Their follow-up laboratory study replicated this effect, but only for video games that were both challenging and enjoyable (ie, when game difficulty was set to “hard” instead of “easy”). Similarly, poor self-control in combination with unstructured television viewing has been associated with delayed bedtimes in adults, a finding supported in a follow-up study.⁹⁴ This emerging research suggests that identifying predisposing individual traits and characteristics of the technology use helps to discern youth whose technology habits are most likely to delay sleep onset.

INTERVENTIONS, POLICIES, AND STRATEGIES DESIGNED TO IMPROVE CHILD AND ADOLESCENT SLEEP

The American Academy of Pediatrics issued a Statement of Endorsement in support of the American Academy of Sleep Medicine guidelines, which recommends that

screen-based devices not be allowed in children's bedrooms and be turned off 30 minutes before bedtime.⁹⁵ These are commonly suggested approaches, but modifying media content may be another effective means of protecting sleep. A randomized trial found that a harm reduction intervention that attempted to change the media exposure of preschool children away from violent and toward educational and prosocial content, significantly decreased odds of sleep problems across follow-up at 6, 12, and 18 months, compared with controls.⁹⁶

Experimental studies seeking to evaluate interventions to improve sleep outcomes in children and adolescents are sparse, and have typically incorporated multifaceted sleep hygiene programs that address multiple elements in addition to screen media before bed. Nevertheless, existing literature suggests that limiting screen time during the 30 to 60 minutes before bedtime may yield modest benefits in terms of "lights out time" as well as sleep quality and duration. A 1-week program, consisting of classes teaching sleep physiology, biological significance, and the consequences of sleep deprivation, resulted in more regular bedtimes and shorter sleep onset latency among 58 adolescents (mean, 16 years of age), but no benefit on sleep quality or daytime sleepiness.⁹⁷ Another investigation assessed the usefulness of the FERRET (Food, Emotions, Routine, Restrict, Environment and Timing) sleep hygiene program among 22 adolescents (mean, 13 years of age). FERRET instructed adolescents to comply with 3 rules for each of the domains of the acronym (eg, restrict—no electronic media at least 30 minutes before bed, no exercise 3 hours before bed, no other activities in bed except for sleep).⁹⁸ This program significantly improved scores on the Adolescent Sleep Hygiene Scale, Pittsburgh Sleep Quality Index, and Sleep Disturbance Scale for Children, although objective sleep duration did not change significantly. A targeted intervention of mobile phone restriction in the hour before bed among 63 adolescents (mean, 16 years of age) resulted in "lights out" 17 minutes earlier and a total sleep time increase of 19 minutes per night.⁹⁹ However, the authors reported a low 26% recruitment rate, highlighting the difficulty of implementing such a regimented screen-limiting schedule in teens. In another study, limiting screen media around bedtime for healthy adolescents with good sleep did not significantly impact sleep outcomes. In a study of high school athletes (mean, 17 years of age) whose baseline consisted of sufficient sleep (eg, total sleep time, 7:49 weekdays, 9:11 weekends), a strict "no electronic media" rule after 10:00 PM resulted in no benefit in terms of total sleep time, sleep quality, or daytime functioning.¹⁰⁰ In sum, limited existing research suggests that sleep hygiene interventions may be practically challenging to achieve but yield benefits to those children and adolescents with insufficient sleep. Larger, multisite studies are urgently needed given the increasing intrusion of evening screen media use in the lives of modern youth, including gaming, social media, and homework.

CLINICIAN'S PERSPECTIVE

Clinicians can help families to improve their sleep health and screen media habits by encouraging parenting marked by high levels of warmth and support, as well as limits that are clearly communicated, consistently applied appropriate to the child's behavior and context, and allow for developmentally appropriate autonomy (ie, an authoritative parenting style^{101,102}; **Box 1**). All parents should begin instilling family bedtime routines and healthy sleep habits early in life, and adjust these routines as youth mature (**Box 2**). If the youth health behaviors and habits become part of the child's own daily routine, she or he will be better able to take charge of her or his own sleep health behaviors when this becomes appropriate in later years. For younger children, these routines mean establishing household rules related to screens and

Box 1**Sleep-friendly screen behavior recommendations for clinicians and educators**

- Talk with families about the importance of adequate sleep.
- Recommend building healthy sleep habits starting as young as possible.
- Teach families about the negative effects of evening use of light-emitting screens on sleep.
- Encourage regular bedtimes that allow adequate time for sleep, and regular bedtime routines in the hour before bed, consisting of calming activities and avoidance of screen media.
- Advise families to restrict all screen devices from bedrooms, including TVs, video games, computers, tablets, and cell phones. Encourage parents to be good role models by following these rules themselves.
- Consider insufficient sleep as a contributing factor for youth exhibiting mood, academic, or behavioral problems.
- Inspire children of all ages to develop autonomy and self-regulatory skills to maintain healthy screen media habits.

Adapted from Lemola S, Perkinson-Gloor N, Brand S, et al. Adolescents' electronic media use at night, sleep disturbance, and depressive symptoms in the smartphone age. *J Youth Adolesc* 2015;44(2):405–18; and Mindell JA, Owens JA. *A clinical guide to pediatric sleep: diagnosis and management of sleep problems*, 2nd edition. Philadelphia: Lippincott Williams & Wilkins; 2015; with permission.

sleep early on. For older children and adolescents, they involve open conversations about the core reasons for behaviors. This proactive and engaged parenting style promotes cooperation and parent–child shared goals for children's health and well-being, and aims to help children develop self-regulation skills, and eventually increasing autonomy to govern their own behavior.

Examples from health psychology provide accessible guidance. A teen with a healthy diet does not typically result from the rigid imposition of a such a diet after a lifetime of unhealthy foods, but with a healthy and balanced eating habits instilled from a young age. A healthy “screen media diet” may be a resonant and useful concept for many children, parents, educators, and clinicians. It is difficult for parents who smoke to mandate a no smoking policy. Role modeling and a health risk prevention approach is far more likely to be effective than a secondary prevention approach (eg, smoking cessation). Successful parenting ideally starts early and sets appropriate norms and boundaries while maintaining parental warmth, rather than holding off on setting boundaries until a problem develops requiring discipline. Encouraging the development of age-appropriate autonomy will help teenagers to develop a sense of responsibility for maintaining a healthy lifestyle.

For teenagers presenting with excessive screen time, a few clinical pearls may help families to follow the recommendations laid out in **Boxes 1** and **2**. First, one must always focus on the chief complaint, that is, what brought the patient to seek help in the first place. In cases of excessive nighttime screen media use, children and adolescents are often seen in the clinician's office for poor academic performance or lack of concentration in school. Upon taking a careful history, the clinician often discovers a significant lack of sleep, often attributable to patients staying up late while using mobile devices, computers, or video games. Clinicians must discover the underlying factors (eg, social or family stress) that drive the patient to use the screens in the late hours. Addressing such factors directly may be essential to motivating families to achieve healthier screen habits.

Box 2**Sleep-friendly screen behavior recommendations for parents**

- Establish screen media habits for your children that enable healthy sleep.^{3,7,111}
 - Plan a bedtime that allows for adequate sleep.
 - Avoid screen media in the hour before bedtime and at nighttime.
 - Replace evening screen time with calm activities for your children (reading, coloring, conversation, etc).
 - Keep all screen devices (televisions, video games, computers, tablets, and smartphones) out of bedrooms.
 - Avoid passive background media: Children may be affected by screen media even when they are not actively engaged.
 - Content matters: Avoid violent and/or scary media, which can negatively affect your children's sleep.
 - Family rules and routines are most effective when applied to all children in the household.
- Establish other healthy sleep practices for your child and yourself.^{112–114}
 - Set and abide by regular bedtimes every day, including weekends, allowing the child sufficient sleep duration for his or her age.¹⁰
 - Bedtime should follow a predictable routine (eg, brush teeth, read a story, lights out).
 - Bedrooms should be cool (65°F–70°F), comfortable, dark, and quiet.
 - Avoid evening intake of chocolate or beverages that interfere with sleep (soda, tea, coffee, energy drinks).
 - Include physical exercise into the daily routine, and spend time outdoors during sunlight hours when possible.
- Be a healthy sleep role model by prioritizing your own sleep.
 - Improve your own sleep-related behaviors (eg, reduce screen time before bedtime, establish a regular bedtime)¹¹³ to improve your health and well-being.¹¹⁵
 - Turn off electronic media devices in the evening throughout the household and charge all mobile devices in a central location outside bedrooms.
 - Parents who are overtired are less well able to parent effectively, including being proactive in orchestrating child routines and dealing effectively and calmly with daily hassles that are part of everyday life.
- Parent your child with clear communication, awareness, and fair, consistent, developmentally appropriate rules.¹¹⁶
 - Talk with your child or teen early and often about the importance of adequate sleep for optimal health.
 - Be aware of how much time your child or teen spends engaging in screen media, including before and after bedtime.⁷⁵
 - Pay attention to your child or teen's mood and behavior at home, and discuss concerns you may have. Mood impairment is often caused by inadequate sleep.
 - Establish and enforce appropriate media and sleep rules for your children as early in their lives as possible; consistently point out after-effects of failing to follow those rules (eg, being tired and cranky the next day after playing games too late) to develop your child's understanding of the effects of inadequate sleep.
 - Work with teens to jointly develop healthy sleep routines that also allow them to meet obligations (eg, homework or sports) and are consistent with these guidelines. Developing autonomy and ability to self-regulate is important for teens, as is consistency.

The clinician should then work with families to reduce evening and nighttime screen media use via structural and behavioral modifications to improve sleep health (see **Box 2**). Many adolescents are reluctant to change their health behaviors,¹⁰³ yet providing them with education about the positive benefits of sufficient sleep as well as the negative consequences of poor sleep can help to motivate such change.^{104–108} For example, pointing out that a change in sleep habits may improve concentration, daytime alertness, and academic performance may motivate youth to modify nighttime habits. Discussing the negative repercussions of inadequate sleep on qualities

that are important to the patient, such as body weight or athletic performance, may likewise motivate the patient to reduce nighttime screen media use and establish other healthy bedtime habits. Perhaps most important, parents and children should remove screen-based devices from the bedroom to ensure that they are not used at night.

Changing bedtime screen media use habits in our young patients is challenging. However, several tools and strategies increase our chances of success, with the goal of promoting healthy sleep habits that children and adolescents will continue to follow throughout their lives.

RECOMMENDATIONS FOR CLINICIANS, EDUCATORS, AND PARENTS

Based on our current understanding of clinical practice and sleep health research, we have developed sleep-friendly screen behavior recommendations for clinicians and educators (see **Box 1**) and for parents (see **Box 2**) to help minimize the adverse effects of screen-based media on the sleep of children and adolescents.

FUTURE DIRECTIONS

Research indicates that screen-based media represents a threat to the sleep quality of youth, many of whom already have insufficient sleep. However, very few studies demonstrate easy-to-implement and effective interventions. Future research should develop, implement, and evaluate sustainable interventions that minimize the adverse effects of evening screen use on sleep. For example, a means of reducing fear of missing out from social media and other screen-based activities may significantly improve the sleep of adolescents.^{109,110} There is a clear need for more basic, translational, and clinical research into the effects of screen-based media on sleep and related health consequences among children and adolescents, to educate and motivate parents, clinicians, teachers, and youth to foster healthy sleep habits.

ACKNOWLEDGMENTS

The authors are grateful to Guest Editor, Paul Weigle, MD, for his careful review and constructive feedback on this article.

REFERENCES

1. National Sleep Foundation (NSF). Children and sleep. 2004. Available at: <https://sleepfoundation.org/sites/default/files/FINAL%20SOF%202004.pdf>. Accessed June 10, 2017.
2. National Sleep Foundation (NSF). Sleep in America poll: teens and sleep. 2006. Available at: https://sleepfoundation.org/sites/default/files/2006_summary_of_findings.pdf. Accessed June 10, 2017.
3. Hale L, Guan S. Screen time and sleep among school-aged children and adolescents: a systematic literature review. *Sleep Med Rev* 2015;21:50–8.
4. Cappuccio FP, Taggart FM, Kandala NB, et al. Meta-analysis of short sleep duration and obesity in children and adults. *Sleep* 2008;31:619–26.
5. Beebe DW. Cognitive, behavioral, and functional consequences of inadequate sleep in children and adolescents. *Pediatr Clin North Am* 2011;58:649–65.
6. Falbe J, Davison KK, Franckle RL, et al. Sleep duration, restfulness, and screens in the sleep environment. *Pediatrics* 2015;135:e367–75.
7. Buxton OM, Chang AM, Spilsbury JC, et al. Sleep in the modern family: protective family routines for child and adolescent sleep. *Sleep Health* 2015;1:15–27.

8. Hysing M, Pallesen S, Stormark KM, et al. Sleep and use of electronic devices in adolescence: results from a large population-based study. *BMJ Open* 2015;5:e006748.
9. Finch KB, Bernstein SJ, Aguilar MD, et al. The RAND/UCLA appropriateness method user's manual. Santa Monica (CA): RAND; 2001.
10. Hirshkowitz M, Whiton K, Albert SM, et al. National Sleep Foundation's sleep time duration recommendations: methodology and results summary. *Sleep Health* 2015;1:40–3.
11. Paruthi S, Brooks LJ, D'Ambrosio C, et al. Consensus statement of the American Academy of Sleep Medicine on the recommended amount of sleep for healthy children: methodology and discussion. *J Clin Sleep Med* 2016;12:1549–61.
12. Sadeh A, Tikotzky L, Kahn M. Sleep in infancy and childhood: implications for emotional and behavioral difficulties in adolescence and beyond. *Curr Opin Psychiatry* 2014;27:453–9.
13. Sivertsen B, Harvey AG, Reichborn-Kjennerud T, et al. Later emotional and behavioral problems associated with sleep problems in toddlers: a longitudinal study. *JAMA Pediatr* 2015;169:575–82.
14. Gregory AM, O'Connor TG. Sleep problems in childhood: a longitudinal study of developmental change and association with behavioral problems. *J Am Acad Child Adolesc Psychiatry* 2002;41:964–71.
15. Miller AL, Seifer R, Crossin R, et al. Toddler's self-regulation strategies in a challenge context are nap-dependent. *J Sleep Res* 2015;24:279–87.
16. Baum KT, Desai A, Field J, et al. Sleep restriction worsens mood and emotion regulation in adolescents. *J Child Psychol Psychiatry* 2014;55:180–90.
17. Sarchiapone M, Mandelli L, Carli V, et al. Hours of sleep in adolescents and its association with anxiety, emotional concerns, and suicidal ideation. *Sleep Med* 2014;15:248–54.
18. Lovato N, Gradisar M. A meta-analysis and model of the relationship between sleep and depression in adolescents: recommendations for future research and clinical practice. *Sleep Med Rev* 2014;18:521–9.
19. Tesler N, Gerstenberg M, Huber R. Developmental changes in sleep and their relationships to psychiatric illnesses. *Curr Opin Psychiatry* 2013;26:572–9.
20. Williams KE, Berthelsen D, Walker S, et al. A developmental cascade model of behavioral sleep problems and emotional and attentional self-regulation across early childhood. *Behav Sleep Med* 2017;15:1–21.
21. Taveras EM, Rifas-Shiman SL, Oken E, et al. Short sleep duration in infancy and risk of childhood overweight. *Arch Pediatr Adolesc Med* 2008;162:305–11.
22. Taveras EM, Gillman MW, Pena MM, et al. Chronic sleep curtailment and adiposity. *Pediatrics* 2014;133:1013–22.
23. Mullins EN, Miller AL, Cherian SS, et al. Acute sleep restriction increases dietary intake in preschool-age children. *J Sleep Res* 2017;26:48–54.
24. Woo Baidal JA, Locks LM, Cheng ER, et al. Risk factors for childhood obesity in the first 1,000 days: a systematic review. *Am J Prev Med* 2016;50:761–79.
25. Franckle RL, Falbe J, Gortmaker S, et al. Insufficient sleep among elementary and middle school students is linked with elevated soda consumption and other unhealthy dietary behaviors. *Prev Med* 2015;74:36–41.
26. Zero to eight: children's media use in America 2013. San Francisco (CA): Common Sense Media Inc.; 2013.
27. The common sense census: media use by tweens and teens. San Francisco (CA): Common Sense Media Inc.; 2015.

28. Rideout V, Hamel E. The media family: electronic media in the lives of infants, toddlers, preschoolers and their parents. Menlo Park (CA): Henry J. Kaiser Family Foundation; 2006.
29. Eggermont S, Van den Bulck J. Nodding off or switching off? The use of popular media as a sleep aid in secondary-school children. *J Paediatr Child Health* 2006;42:428–33.
30. Cain N, Gradisar M. Electronic media use and sleep in school-aged children and adolescents: a review. *Sleep Med* 2010;11:735–42.
31. Bartel KA, Gradisar M, Williamson P. Protective and risk factors for adolescent sleep: a meta-analytic review. *Sleep Med Rev* 2015;21:72–85.
32. Carter B, Rees P, Hale L, et al. Association between portable screen-based media device access or use and sleep outcomes: a systematic review and meta-analysis. *JAMA Pediatr* 2016;170:1202–8.
33. Harbard E, Allen NB, Trinder J, et al. What's keeping teenagers up? prebedtime behaviors and actigraphy-assessed sleep over school and vacation. *J Adolesc Health* 2016;58:426–32.
34. Fobian AD, Avis K, Schwebel DC. Impact of media use on adolescent sleep efficiency. *J Dev Behav Pediatr* 2016;37:9–14.
35. Muller D, Signal L, Elder D, et al. Environmental and behavioural factors associated with school children's sleep in Aotearoa/New Zealand. *J Paediatr Child Health* 2017;53:68–74.
36. Exelmans L, Van den Bulck J. Sleep quality is negatively related to video gaming volume in adults. *J Sleep Res* 2015;24:189–96.
37. Vijakkhana N, Wilaisakditipakorn T, Ruedeekhajorn K, et al. Evening media exposure reduces night-time sleep. *Acta Paediatr* 2015;104:306–12.
38. Sijsma A, Koller M, Sauer PJ, et al. Television, sleep, outdoor play and BMI in young children: the GECKO Drenthe cohort. *Eur J Pediatr* 2015;174:631–9.
39. Chaput JP, Leduc G, Boyer C, et al. Electronic screens in children's bedrooms and adiposity, physical activity and sleep: do the number and type of electronic devices matter? *Can J Public Health* 2014;105:e273–9.
40. Al-Hazzaa HM, Al-Sobayel HI, Abahussain NA, et al. Association of dietary habits with levels of physical activity and screen time among adolescents living in Saudi Arabia. *J Hum Nutr Diet* 2014;27(Suppl 2):204–13.
41. Jiang X, Hardy LL, Baur LA, et al. Sleep duration, schedule and quality among urban Chinese children and adolescents: associations with routine after-school activities. *PLoS One* 2015;10:e0115326.
42. Yland J, Guan S, Emanuele E, et al. Interactive vs passive screen time and night-time sleep duration among school-aged children. *Sleep Health* 2015;1:191–6.
43. King DL, Delfabbro PH, Zwaans T, et al. Sleep interference effects of pathological electronic media use during adolescence. *Int J Ment Health* 2013;1.
44. Nuutinen T, Ray C, Roos E. Do computer use, TV viewing, and the presence of the media in the bedroom predict school-aged children's sleep habits in a longitudinal study? *BMC Public Health* 2013;13:684.
45. Oka Y, Suzuki S, Inoue Y. Bedtime activities, sleep environment, and sleep/wake patterns of Japanese elementary school children. *Behav Sleep Med* 2008;6:220–33.
46. Ivarsson M, Anderson M, Akerstedt T, et al. The effect of violent and nonviolent video games on heart rate variability, sleep, and emotions in adolescents with different violent gaming habits. *Psychosom Med* 2013;75:390–6.
47. Munezawa T, Kaneita Y, Osaki Y, et al. The association between use of mobile phones after lights out and sleep disturbances among Japanese adolescents: a nationwide cross-sectional survey. *Sleep* 2011;34:1013–20.

48. Alexandru G, Michikazu S, Shimako H, et al. Epidemiological aspects of self-reported sleep onset latency in Japanese junior high school children. *J Sleep Res* 2006;15:266–75.
49. Wallenius M, Punamäki RL, Rimpelä A. Digital game playing and direct and indirect aggression in early adolescence: the roles of age, social intelligence, and parent-child communication. *J Youth Adolescence* 2007;36:325–36.
50. Lemola S, Brand S, Vogler N, et al. Habitual computer game playing at night is related to depressive symptoms. *Personality Individ Differ* 2011;51:117–22.
51. Owens J, Maxim R, McGuinn M, et al. Television-viewing habits and sleep disturbance in school children. *Pediatrics* 1999;104:e27.
52. Van den Bulck J. Television viewing, computer game playing, and Internet use and self-reported time to bed and time out of bed in secondary-school children. *Sleep* 2004;27:101–4.
53. Garrison MM, Liekweg K, Christakis DA. Media use and child sleep: the impact of content, timing, and environment. *Pediatrics* 2011;128:29–35.
54. Mazurek MO, Engelhardt CR, Hilgard J, et al. Bedtime electronic media use and sleep in children with autism spectrum disorder. *J Dev Behav Pediatr* 2016;37:525–31.
55. Johnson JG, Cohen P, Kasen S, et al. Association between television viewing and sleep problems during adolescence and early adulthood. *Arch Pediatr Adolesc Med* 2004;158:562–8.
56. Li S, Jin X, Wu S, et al. The impact of media use on sleep patterns and sleep disorders among school-aged children in China. *Sleep* 2007;30:361–7.
57. Thompson DA, Christakis DA. The association between television viewing and irregular sleep schedules among children less than 3 years of age. *Pediatrics* 2005;116:851–6.
58. Adam EK, Snell EK, Pendry P. Sleep timing and quantity in ecological and family context: a nationally representative time-diary study. *J Fam Psychol* 2007;21:4–19.
59. Dworak M, Schierl T, Bruns T, et al. Impact of singular excessive computer game and television exposure on sleep patterns and memory performance of school-aged children. *Pediatrics* 2007;120:978–85.
60. Higuchi S, Motohashi Y, Liu Y, et al. Effects of playing a computer game using a bright display on presleep physiological variables, sleep latency, slow wave sleep and REM sleep. *J Sleep Res* 2005;14:267–73.
61. Yen CF, Ko CH, Yen JY, et al. The multidimensional correlates associated with short nocturnal sleep duration and subjective insomnia among Taiwanese adolescents. *Sleep* 2008;31:1515–25.
62. Choi K, Son H, Park M, et al. Internet overuse and excessive daytime sleepiness in adolescents. *Psychiatry Clin Neurosci* 2009;63:455–62.
63. Punamaki RL, Wallenius M, Nygard CH, et al. Use of information and communication technology (ICT) and perceived health in adolescence: the role of sleeping habits and waking-time tiredness. *J Adolesc* 2007;30:569–85.
64. Mesquita G, Reimao R. Nightly use of computer by adolescents: its effect on quality of sleep. *Arq Neuropsiquiatr* 2007;65:428–32.
65. Arora T, Broglio E, Thomas GN, et al. Associations between specific technologies and adolescent sleep quantity, sleep quality, and parasomnias. *Sleep Med* 2014;15:240–7.
66. Arora T, Hussain S, Hubert Lam KB, et al. Exploring the complex pathways among specific types of technology, self-reported sleep duration and body mass index in UK adolescents. *Int J Obes (Lond)* 2013;37:1254–60.

67. Weaver E, Gradisar M, Dohnt H, et al. The effect of presleep video-game playing on adolescent sleep. *J Clin Sleep Med* 2010;6:184–9.
68. Lemola S, Perkinson-Gloor N, Brand S, et al. Adolescents' electronic media use at night, sleep disturbance, and depressive symptoms in the smartphone age. *J Youth Adolesc* 2015;44:405–18.
69. Chahal H, Fung C, Kuhle S, et al. Availability and night-time use of electronic entertainment and communication devices are associated with short sleep duration and obesity among Canadian children. *Pediatr Obes* 2013;8:42–51.
70. Gradisar M, Wolfson AR, Harvey AG, et al. The sleep and technology use of Americans: findings from the National Sleep Foundation's 2011 Sleep in America poll. *J Clin Sleep Med* 2013;9:1291–9.
71. King DL, Gradisar M, Drummond A, et al. The impact of prolonged violent video-gaming on adolescent sleep: an experimental study. *J Sleep Res* 2013;22:137–43.
72. Heath M, Sutherland C, Bartel K, et al. Does one hour of bright or short-wavelength filtered tablet screenlight have a meaningful effect on adolescents' pre-bedtime alertness, sleep, and daytime functioning? *Chronobiol Int* 2014;31:496–505.
73. Cajochen C, Frey S, Anders D, et al. Evening exposure to a light-emitting diodes (LED)-backlit computer screen affects circadian physiology and cognitive performance. *J Appl Physiol (1985)* 2011;110:1432–8.
74. van der Lely S, Frey S, Garbazza C, et al. Blue blocker glasses as a countermeasure for alerting effects of evening light-emitting diode screen exposure in male teenagers. *J Adolesc Health* 2015;56:113–9.
75. Chang AM, Aeschbach D, Duffy JF, et al. Evening use of light-emitting eReaders negatively affects sleep, circadian timing, and next-morning alertness. *Proc Natl Acad Sci U S A* 2015;112:1232–7.
76. Wood B, Rea MS, Plitnick B, et al. Light level and duration of exposure determine the impact of self-luminous tablets on melatonin suppression. *Appl Ergon* 2013;44:237–40.
77. Bruni O, Sette S, Fontanesi L, et al. Technology use and sleep quality in preadolescence and adolescence. *J Clin Sleep Med* 2015;11:1433–41.
78. Reynolds CM, Gradisar M, Kar K, et al. Adolescents who perceive fewer consequences of risk-taking choose to switch off games later at night. *Acta Paediatr* 2015;104:e222–7.
79. van der Vijgh B, Beun RJ, Van Rood M, et al. Meta-analysis of digital game and study characteristics eliciting physiological stress responses. *Psychophysiology* 2015;52:1080–98.
80. Barlett CP, Rodeheffer C. Effects of realism on extended violent and nonviolent video game play on aggressive thoughts, feelings, and physiological arousal. *Aggress Behav* 2009;35:213–24.
81. Lin TC. Effects of gender and game type on autonomic nervous system physiological parameters in long-hour online game players. *Cyberpsychol Behav Soc Netw* 2013;16:820–7.
82. Ivarsson M, Anderson M, Akerstedt T, et al. Playing a violent television game affects heart rate variability. *Acta Paediatr* 2009;98:166–72.
83. Higuchi S, Motohashi Y, Liu Y, et al. Effects of VDT tasks with a bright display at night on melatonin, core temperature, heart rate, and sleepiness. *J Appl Physiol (1985)* 2003;94:1773–6.
84. Lockley SW, Brainard GC, Czeisler CA. High sensitivity of the human circadian melatonin rhythm to resetting by short wavelength light. *J Clin Endocrinol Metab* 2003;88:4502–5.

85. Turner PL, Mainster MA. Circadian photoreception: ageing and the eye's important role in systemic health. *Br J Ophthalmol* 2008;92:1439–44.
86. Higuchi S, Nagafuchi Y, Lee SI, et al. Influence of light at night on melatonin suppression in children. *J Clin Endocrinol Metab* 2014;99:3298–303.
87. Crowley SJ, Cain SW, Burns AC, et al. Increased sensitivity of the circadian system to light in early/mid-puberty. *J Clin Endocrinol Metab* 2015;100:4067–73.
88. Association Psychiatric Association. Diagnostic and statistical manual of mental disorders (5th edition). Washington, DC: American Psychiatric Publishing; 2013.
89. Goldsmith TD, Shapira NA. Problematic internet use. In: Hollander E, Stein DJ, editors. *Clinical manual of impulse-control disorders*. 1st edition. Washington, DC: American Psychiatric Publishing; 2005. p. 291–308.
90. Lam LT. Internet gaming addiction, problematic use of the internet, and sleep problems: a systematic review. *Curr Psychiatry Rep* 2014;16:444.
91. Achab S, Nicolier M, Mauny F, et al. Massively multiplayer online role-playing games: comparing characteristics of addict vs non-addict online recruited gamers in a French adult population. *BMC Psychiatry* 2011;11:144.
92. Rehbein F, Kleimann M, Mossle T. Prevalence and risk factors of video game dependency in adolescence: results of a German nationwide survey. *Cyberpsychol Behav Soc Netw* 2010;13:269–77.
93. Smith LJ, Gradisar M, King DL, et al. Intrinsic and extrinsic predictors of video-gaming behaviour and adolescent bedtimes: the relationship between flow states, self-perceived risk-taking, device accessibility, parental regulation of media and bedtime. *Sleep Med* 2017;30:64–70.
94. Exelmans, L, VD, BJ. Ego depletion both increases and decreases time to bed: a dual pathway model. Presented at the 23rd Congress of the European Sleep Research Society, Bologna (Italy), September 13–16, 2016.
95. American Academy of Pediatrics Supports Childhood Sleep Guidelines. American Academy of Pediatrics (2016).
96. Garrison MM, Christakis DA. The impact of a healthy media use intervention on sleep in preschool children. *Pediatrics* 2012;130:492–9.
97. de Sousa IC, Araujo JF, de Azevedo CVM, et al. The effect of a sleep hygiene education program on the sleep–wake cycle of Brazilian adolescent students. *Sleep Biol Rhythms* 2007;5:251–8.
98. Tan E, Healey D, Gray AR, et al. Sleep hygiene intervention for youth aged 10 to 18 years with problematic sleep: a before–after pilot study. *BMC Pediatr* 2012;12:189.
99. Bartel K, Gradisar M. An adolescent's worst nightmare? Altering pre-bedtime phone use to achieve better sleep health. *J Sleep Res* 2016;25:141.
100. Harris A, Gundersen H, Mørk-Andreassen P, et al. Restricted use of electronic media, sleep, performance, and mood in high school athletes—a randomized trial. *Sleep Health* 2015;1:314–21.
101. Klein HA, Ballantine J. For parents particularly: raising competent kids: the authoritative parenting style. *Child Education* 2012;79(1):46–7.
102. Grusec JE. A history of research on parenting strategies and children's internalization of values. In: Grusec JE, Kuczynski L, editors. *Parenting and children's internalization of values*. Hoboken (NJ): Wiley; 1997. p. 3–22.
103. Gayes LA, Steele RG. A meta-analysis of motivational interviewing interventions for pediatric health behavior change. *J Consult Clin Psychol* 2014;82:521–35.
104. Prochaska JO, DiClemente CC. Stages and processes of self-change of smoking: toward an integrative model of change. *J Consult Clin Psychol* 1983;51:390–5.

105. Cain N, Gradisar M, Moseley L. A motivational school-based intervention for adolescent sleep problems. *Sleep Med* 2011;12:246–51.
106. Bonnar D, Gradisar M, Moseley L, et al. Evaluation of novel school-based interventions for adolescent sleep problems: does parental involvement and bright light improve outcomes? *Sleep Health* 2015;1:66–74.
107. Cassoff J, Knauper B, Michaelsen S, et al. School-based sleep promotion programs: effectiveness, feasibility and insights for future research. *Sleep Med Rev* 2013;17:207–14.
108. Gradisar M, Smits MG, Bjorvatn B. Assessment and treatment of delayed sleep phase disorder in adolescents: recent innovations and cautions. *Sleep Med Clin* 2014;9:199–210.
109. Beyens I, Frison E, Eggermont S. “I don’t want to miss a thing”: adolescents’ fear of missing out and its relationship to adolescents’ social needs, Facebook use, and Facebook related stress. *Comput Hum Behav* 2016;64:1–8.
110. Elhai JD, Levine JC, Dvorak RD, et al. Fear of missing out, need for touch, anxiety and depression are related to problematic smartphone use. *Comput Hum Behav* 2016;63:509–16.
111. National Sleep Foundation (NSF). Electronics in the bedroom: Why it’s necessary to turn off before you tuck in. 2017. Available at: <https://sleepfoundation.org/ask-the-expert/electronics-the-bedroom>. Accessed June 10, 2017.
112. Mindell J, Owens J. *A clinical guide to pediatric sleep: diagnosis and management of sleep problems*. Philadelphia: Lippincott Williams & Wilkins; 2015.
113. National Sleep Foundation (NSF). Back to school sleep tips. 2017. Available at: <http://sleepfoundation.org/sleep-news/back-school-sleep-tips-0>. Accessed June 10, 2017.
114. National Sleep Foundation (NSF). Healthy sleep tips. 2017. Available at: <https://sleepfoundation.org/sleep-tools-tips/healthy-sleep-tips>. Accessed June 10, 2017.
115. Watson NF, Badr MS, Belenky G, et al. Recommended amount of sleep for a healthy adult: a joint consensus statement of the American Academy of Sleep Medicine and sleep research society. *Sleep* 2015;38:843–4.
116. Positive Parenting Practices. Center for Disease Control and Prevention. 2016. Available at: <http://www.cdc.gov/healthyyouth/protective/positiveparenting/monitoring.htm>. Accessed June 10, 2017.